



# Role of Neuraminidase in Influenza A(H7N9) Virus Receptor Binding

Donald J. Benton,<sup>a</sup> Stephen A. Wharton,<sup>a</sup> Stephen R. Martin,<sup>b</sup> John W. McCauley<sup>a</sup>

Worldwide Influenza Centre<sup>a</sup> and Structural Biology Science Technology Platform,<sup>b</sup> Francis Crick Institute, London, United Kingdom<sup>b</sup>

**ABSTRACT** Influenza A(H7N9) viruses have caused a large number of zoonotic infections since their emergence in 2013. They remain a public health concern due to the repeated high levels of infection with these viruses and their perceived pandemic potential. A major factor that determines influenza A virus fitness and therefore transmissibility is the interaction of the surface glycoproteins hemagglutinin (HA) and neuraminidase (NA) with the cell surface receptor sialic acid. Typically, the HA is responsible for binding to the sialic acid to allow virus internalization and the NA is a sialidase responsible for cleaving sialic acid to aid virus spread and release. N9 NA has previously been shown to have receptor binding properties mediated by a sialic acid binding site, termed the hemadsorption (Hb) site, which is discrete from the enzymatically active sialidase site. This study investigated the N9 NA from a zoonotic H7N9 virus strain in order to determine its possible role in virus receptor binding. We demonstrate that this N9 NA has an active Hb site which binds to sialic acid, which enhances overall virus binding to sialic acid receptor analogues. We also show that the N9 NA can also contribute to receptor binding due to unusual kinetic characteristics of the sialidase site which specifically enhance binding to human-like  $\alpha,6$ -linked sialic acid receptors.

**IMPORTANCE** The interaction of influenza A virus glycoproteins with cell surface receptors is a major determinant of infectivity and therefore transmissibility. Understanding these interactions is important for understanding which factors are necessary to determine pandemic potential. Influenza A viruses generally mediate binding to cell surface sialic acid receptors via the hemagglutinin (HA) glycoprotein, with the neuraminidase (NA) glycoprotein being responsible for cleaving the receptor to allow virus release. Previous studies showed that the NA proteins of the N9 subtype can bind sialic acid via a separate binding site distinct from the sialidase active site. This study demonstrates for purified protein and virus that the NA of the zoonotic H7N9 viruses has a binding capacity via both the secondary binding site and unusual kinetic properties of the sialidase site which promote receptor binding via this site and which enhance binding to human-like receptors. This could have implications for understanding human-to-human transmission of these viruses.

**KEYWORDS** hemagglutinin, influenza A virus, neuraminidase, receptor analogues, receptor binding, biophysics, enzyme kinetics

Influenza A(H7N9) viruses first emerged as a human infection in March 2013. These viruses have been responsible for >1,250 confirmed infections, with a case fatality rate in the region of 25% (1). These infections have occurred in five separate waves, one each year since 2013. The infection of humans with these viruses is zoonotic, with very limited evidence of human-to-human transmission (2). Due to the ability of these viruses to infect humans and cause severe disease, their transmission characteristics are therefore of interest in order to assess their pandemic potential.

Received 5 December 2016 Accepted 14 March 2017

Accepted manuscript posted online 29 March 2017

**Citation** Benton DJ, Wharton SA, Martin SR, McCauley JW. 2017. Role of neuraminidase in influenza A(H7N9) virus receptor binding. *J Virol* 91:e02293-16. <https://doi.org/10.1128/JVI.02293-16>.

**Editor** Bryan R. G. Williams, Hudson Institute of Medical Research

**Copyright** © 2017 Benton et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to John W. McCauley, [john.mccauley@crick.ac.uk](mailto:john.mccauley@crick.ac.uk).


Major factors determining virus fitness and therefore the transmissibility of influenza A virus are the characteristics of the two surface glycoproteins, hemagglutinin (HA) and neuraminidase (NA), which control the interaction of the virus with the cell surface. These two proteins have antagonistic activities, with the HA being responsible for binding to the receptor sialic acid on the surface of cells, whereas the NA is a sialidase responsible for releasing sialic acid from glycoprotein and glycolipid sialoconjugates, to which virus may be bound, to aid virus release. A large number of previous studies have investigated the interdependence of these two activities (reviewed in reference 3).

NA from certain virus subtypes, particularly N9, have been shown to have a secondary sialic acid binding site, the hemadsorption (Hb) site, in addition to the catalytic sialidase site (4, 5). The presence of sialic acid binding via this Hb site has been shown to enhance the catalytic rate for cleavage of large multimeric substrates, such as heavily sialylated glycoproteins (5). The role of this Hb site in virus receptor binding remains unclear. Certain N2 NAs have also been found to have receptor binding properties, which occur via the sialidase site of the NA due to the substitution D151G (6). These D151G mutant NAs were found to have receptor binding properties due to their unusual enzymatic properties, with a low  $K_m$  and a low  $k_{cat}$  indicating that the NA binds the substrate with greater strength but with lower rates of enzymatic cleavage, meaning that the NA acquires more of a receptor binding role (7).

The receptor binding properties of a zoonotic H7N9 virus have previously been characterized by biolayer interferometry (BLI) (8). This and other studies (9–15) indicate that the H7 HA has an overall preference for binding to avian-like  $\alpha 2,3$ -linked sialic acid receptors, but the viruses also show considerable binding to the human-like  $\alpha 2,6$ -linked sialic acid receptors. It is unclear, however, how these binding characteristics affect the ability of these viruses to result in zoonotic infection but not transmit between humans. This study investigates the receptor binding and enzymatic characteristics of the N9 NA from a prototype H7N9 virus with an aim to elucidate a possible role of the NA in affecting virus binding properties. We present biophysical data, obtained using previously characterized techniques (16), that show that the N9 NA can bind to sialic acid via the secondary Hb site and that the sialidase site enhances virus binding to sialic acid, particularly to human-like  $\alpha 2,6$ -linked receptors. The enhancement of binding seen with this particular NA has implications for understanding the fitness and therefore the transmissibility of these viruses.

## RESULTS AND DISCUSSION


**Binding properties of recombinant N9 protein.** Experiments were carried out to determine whether the N9 NA of the H7N9 viruses infecting humans has receptor binding properties. The N9 NA of A/Anhui/1/2013 (Anhui13) was expressed in insect cells as the wild type (WT) and as a previously characterized S367N mutant, which is known to abolish sialic acid binding via the Hb site (5). These expressed proteins were attached to His tag binding magnetic beads and used to measure the ability of the protein to capture turkey red blood cells (TRBCs). NA-coated beads were incubated with TRBCs, and the beads were rapidly pelleted using a magnet. The blood remaining in the supernatant either was left to settle in a microtiter plate (Fig. 1A) or was lysed, and the relative hemoglobin concentration was determined by absorbance spectroscopy (Fig. 1B). These assays demonstrated the ability of WT NA to bind to TRBCs, thus demonstrating its sialic acid binding properties. This binding was abolished when the substitution S367N was introduced, demonstrating that the majority of the binding seen is mediated through the Hb site. Similar levels of binding by the N9 NA were seen in the presence and absence of the sialidase site inhibitor oseltamivir carboxylate, ruling out binding of this inhibitor to the Hb site. There was, however, enhanced capture of TRBCs, indicated by a reduced level of residual TRBCs following capture by the NA-coated beads, when the sialidase site of the mutant NA was not inhibited for both the WT and S367N proteins (Fig. 1B), suggesting a possible contribution of this enzyme active site to receptor binding.



**FIG 1** Red blood cell capture assay by WT (Hb<sup>+</sup>) and S367N (Hb<sup>-</sup>) N9 NAs. Insect cell-expressed N9 NA was attached to His tag binding magnetic beads. The beads were incubated with turkey red blood cells (TRBCs) and rapidly pelleted with a magnet. The supernatant was removed and left to settle in a microtiter plate (A), or TRBCs were lysed with SDS and the released hemoglobin was quantified by measuring the  $A_{540}$  (B). The results of control experiments without protein attached to beads (no protein) and without incubation of beads with TRBCs (no beads) are also shown. Absorbance measurements were normalized to those for the no-protein control and are shown as the means from three independent measurements, with error bars showing standard deviations from the mean. When present, oseltamivir carboxylate was added at a concentration of 100  $\mu$ M. \*\*,  $P < 0.01$ .

**Virus binding properties.** A number of different viruses which contained the NA from Anhui13 in both the wild-type form (which bound via the Hb site [Hb<sup>+</sup> NA]) and the S367N mutant form (which lacked binding via the Hb site [Hb<sup>-</sup> NA]) were generated by reverse genetics to examine the contribution of the Hb site to receptor binding. The viruses generated consisted of H7N9 viruses with the HA from Anhui13, H1N9 viruses with the HA from A/Puerto Rico/8/34 (PR8), and H3N9 viruses with the HA derived from the recent cell culture-propagated cultivar of A/Victoria/361/2012 (Vic361) H3N2 virus (17).

The equilibrium receptor binding characteristics of these viruses were determined by biolayer interferometry (BLI), measuring virus binding as a function of relative sugar loading (RSL) in the presence of NA inhibitors, as previously described (18). Figure 2 shows the results for H7N9 and H1N9 virus binding to sialoglycopolymers bearing the human-like  $\alpha$ 2,6-sialyl-*N*-acetylglucosamine (6SLN) and avian-like  $\alpha$ 2,3-sialyl-*N*-acetylglucosamine (3SLN) receptor analogues. The observed binding of the H7N9 virus was similar to previously reported BLI data for binding of the wild-type virus (8), with an overall preference for binding to the avian-like receptor 3SLN and marginally weaker binding to the human receptor analogue 6SLN. The H7N9 and H1N9 viruses showed an enhancement of binding to both 6SLN and 3SLN when the Hb<sup>+</sup> NA was present, giving an estimated decrease in the relative  $K_{d(\text{virus})}$  (dissociation constant for the virus) of  $\sim$ 2-fold. The H1N9 virus with Hb<sup>-</sup> NA (H1N9 Hb<sup>-</sup> virus) showed levels of binding similar to those of the full 8-segment PR8 virus (Fig. 2B), indicating that the mutation (S367N) reduces binding to a level which is very close to that of the control virus, PR8, without Hb binding in the NA. Both H7N9 and H1N9 viruses showed similar




**FIG 2** Bi-layer interferometry curves of influenza A virus (100 pM) binding to the receptor analogues  $\alpha$ 2,6-sialyl-*N*-acetylglucosamine (6SLN) and  $\alpha$ 2,3-sialyl-*N*-acetylglucosamine (3SLN). The binding of H7N9 (A) and H1N9 (B) viruses to both wild-type (Hb<sup>+</sup>) and S367N mutant (Hb<sup>-</sup>) N9 NAs was measured. The plotted data are the fractional saturation of virus binding as a function of sugar loading.

increases in binding to both the  $\alpha$ 2,6- and  $\alpha$ 2,3-linked receptors when the Hb<sup>+</sup> NA was present. This indicates that there is no receptor linkage preference associated with the presence of the Hb<sup>+</sup> NA.

H3N9 viruses with the HA from Vic361 were constructed in order to assess whether the weak binding to human-like receptor analogues of these recent H3N2 viruses (17, 18) could be enhanced by the addition of an Hb<sup>+</sup> NA. Binding of these viruses is weak, and only binding to 6SLN is observed; consequently, measurements had to be made at 50 times the normal virus concentration (5 nM). At this concentration, the presence of the Hb<sup>+</sup> NA gave an increase in binding to 6SLN (Fig. 3A and B), but no binding to any other receptors tested was observed. This increase in binding amplitude to a saturation level of  $\sim$ 0.75 nm is, however, small compared to the binding amplitude in similar experiments for H1N9 and H7N9 viruses, which give signals of  $>4$  nm at saturation. This apparent preference for 6SLN is likely not due to a specificity of the N9 Hb site for 6SLN, but rather, the binding via the Hb site complements the weak binding of the HA, which is to 6SLN. It is notable that viruses bearing the HA from recent H3N2 viruses normally require MDCK-SIAT cells for efficient replication (6, 19); these cells have enhanced expression of  $\alpha$ 2,6-linked cell surface receptors (20). The propagation characteristics of Hb<sup>+</sup> and Hb<sup>-</sup> H3N9 viruses were similar in MDCK-SIAT cells (Fig. 3C); however, the Hb<sup>+</sup> virus had enhanced replication in MDCK cells, indicating that the Hb site can enhance binding to the insufficient receptors present on MDCK cells.

**HA/NA balance characteristics.** The balance between the activities of the HA and NA was measured using an approach developed previously (16). In these experiments, the binding of viruses was measured both in the presence and in the absence of NA inhibitors. Figure 4 shows the binding of H7N9 viruses with Hb<sup>+</sup> and Hb<sup>-</sup> NA binding to the human-like 6SLN and the avian-like 3SLN receptors. The binding of the Hb<sup>+</sup> and Hb<sup>-</sup> H7N9 viruses to 3SLN showed behavior (Fig. 4B and D) similar to that measured previously with the H3N2 virus X-31 (16), with the uninhibited virus showing very little




**FIG 3** (A and B) Binding of H3N9 viruses to Hb<sup>+</sup> (A) and Hb<sup>-</sup> (B) NAs determined by biolayer interferometry. Binding to a range of different sugars was measured:  $\alpha$ 2,6- and  $\alpha$ 2,3-sialyl-*N*-acetylglucosamine (6SLN and 3SLN, respectively),  $\alpha$ 2,6- and  $\alpha$ 2,3-sialyllactose (6SL and 3SL, respectively), 3SLN 6' sulfated on GlcNAc (6-Su-3SLN), sialyl-Lewis X (Sia Lex), and sialyl Lewis X 6' sulfated on GlcNAc (6-Su-Sia Lex). The data shown are experimental response traces. (C) Plaque assay of H3N9 viruses with Hb<sup>+</sup> and Hb<sup>-</sup> NAs. The results of assays carried out using both MDCK and MDCK-SIAT cells are shown in duplicate. The plaques shown for MDCK cells used 10 times the concentration of each input virus compared to that used for MDCK-SIAT cells.

binding, presumably due to efficient cleavage of the  $\alpha$ 2,3-linked receptor by the NA. Binding to the human-like receptor 6SLN showed different characteristics. In the absence of NA inhibitors, the H7N9 Hb<sup>+</sup> virus showed an initial enhancement of binding at short times (<500 s) which was not seen with the Hb<sup>-</sup> virus (Fig. 4A and C). This enhancement of binding, when taken at a single time point of 250 s, provided a 23% higher binding signal in the absence of NA inhibitors. This binding enhancement was seen only when the Hb site was present and when the sialidase site was uninhibited. Although it is likely that both the sialidase site and the Hb site contribute to this enhancement of binding, it is not possible to assess the relative importance of the two sites with the data obtained. The binding of the H7N9 Hb<sup>+</sup> virus to 6SLN in the absence of NA inhibitors showed a lower fractional saturation maximum (~0.6) than did that of the Hb<sup>-</sup> virus (~0.8) (Fig. 4A and C), suggesting that the presence of the Hb site could enhance the ability of the NA to cleave multimeric substrates, as previously reported (5).

A similar enhancement of binding to 6SLN was observed when the H1N9 Hb<sup>+</sup> and Hb<sup>-</sup> viruses were compared (Fig. 5), although the enhancement of binding was smaller than that seen for the H7N9 viruses, likely due to the higher affinity of the PR8 HA for 6SLN (Fig. 2B), which would lead to a smaller proportional binding contribution of the NA to these viruses.

**NA kinetics.** The data presented above for virus binding indicate that the N9 NA enhances overall receptor binding in the context of a number of different viruses. This binding is mediated by both the Hb and the sialidase sites of the NA. The binding mediated through the sialidase site appears to preferentially result in binding to the  $\alpha$ 2,6-linked receptor 6SLN. It is, consequently, important to identify the kinetic parameters for different receptors to understand these unusual binding characteristics of the N9 NA.






**FIG 4** Biolayer interferometry curves of binding of H7N9 viruses with wild-type (Hb<sup>+</sup>) and S367N (Hb<sup>-</sup>) NA to the receptor analogues  $\alpha$ 2,6-sialyl-N-acetylglucosamine (6SLN) and  $\alpha$ 2,3-sialyl-N-acetylglucosamine (3SLN). Measurements were made in the presence (solid lines) and absence (dashed lines) of NA inhibitors (Inhib).

Enzyme kinetic parameters were determined for expressed Hb<sup>+</sup> and Hb<sup>-</sup> Anhui13 NAs. The cleavage of the fluorogenic substrate 2'-(4-methylumbelliferyl)- $\alpha$ -D-N-acetylneuraminic acid (MUNANA) was the same for both the Hb<sup>+</sup> and Hb<sup>-</sup> NAs, indicating that there is no influence of the Hb site on the ability of the NA to cleave small monomeric substrates (Table 1).

The cleavage of the monomeric avian-like receptor 3SLN is efficient, with a relatively low  $K_m$  and a high  $k_{cat}$ , giving results similar to those previously determined for the NA of X-31 (Table 1) (16). The characteristics of cleavage for the human-like receptor 6SLN are, however, unusual compared to those previously obtained for the NA of the H3N2 virus X-31. The  $K_m$  was  $\sim$ 2-fold lower ( $4,179 \pm 615 \mu\text{M}$  for Anhui13,  $8,070 \pm 615 \mu\text{M}$  for X-31), indicating stronger substrate binding. The  $k_{cat}$  of the N9 NA was also  $\sim$ 10-fold lower than that of the X-31 NA ( $2.2 \pm 0.2 \text{ s}^{-1}$  for Anhui13,  $18.5 \pm 0.9 \text{ s}^{-1}$  for X-31), indicating a reduced enzymatic turnover (16). The combination of these two parameters shows that the N9 NA can bind to  $\alpha$ 2,6-linked receptors more strongly but cleaves them with a lower efficiency than does X-31, suggesting that the N9 NA has the ability to play a receptor binding role. If one assumes a reasonable value for the association constant ( $k_1$ ) of sialic acid for the NA sialidase site of  $5 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$  (21), the substrate dissociation constant ( $k_{-1}$ ) can be calculated using the  $K_m$  and  $k_{cat}$  values determined for the N9 NA ( $k_{-1} = K_m \cdot k_1 - k_{cat}$ ). Using the calculated  $k_{-1}$  and the experimentally determined  $k_{cat}$ , the probability that bound substrate will be cleaved or will dissociate before undergoing cleavage can be determined by calculating the ratio between  $k_{cat}$  and  $k_{-1}$ . The N9 NA is  $\sim$ 1,000-fold more likely to dissociate rather than cleave 6SLN, whereas the likelihood for 3SLN is  $\sim$ 1.5-fold. The comparable probability values for another NA previously characterized from the H3N2 virus X-31 (16) are an  $\sim$ 200-fold probability of 6SLN dissociation rather than cleavage and an  $\sim$ 1.8-fold probability for 3SLN dissociation rather than cleavage. Therefore, the N9 NA has an  $\sim$ 5-fold enhanced probability of 6SLN dissociation compared to the NA of X-31, and the probability value for 3SLN is similar. This enhanced probability of 6SLN dissociation rather than cleavage



**FIG 5** Bi-layer interferometry curves of H1N9 virus (100 pM) binding to the receptor analogues  $\alpha$ 2,6-sialyl-*N*-acetylglucosamine (6SLN) and  $\alpha$ 2,3-sialyl-*N*-acetylglucosamine (3SLN). Measurements were made in the presence (solid lines) and absence (dashed lines) of NA inhibitors. The binding of viruses to wild-type (Hb<sup>+</sup>) and S367N (Hb<sup>-</sup>) N9 NAs was measured.

coupled with a relatively low  $K_m$  compared to that of the X-31 NA reinforces the hypothesis that substrate binding via the sialidase site is at least partially responsible for the enhancement of the initial binding to 6SLN by H7N9 and H1N9 viruses with the WT Hb<sup>+</sup> NA (Fig. 4 and 5).

Cleavage of the multimeric substrate fetuin by the Hb<sup>+</sup> NA has a  $K_m$  ( $197.5 \pm 37.3 \mu\text{M}$ ) lower than that measured for the Hb<sup>-</sup> NA ( $346.4 \pm 95.6 \mu\text{M}$ ), indicating that the presence of the Hb site increases the affinity of the NA for multimeric substrates. However, the  $k_{\text{cat}}$  values for the Hb<sup>+</sup> and the Hb<sup>-</sup> NAs are similar, indicating that there is no difference in enzyme turnover when the substrate concentration is not limiting. The Hb<sup>-</sup> NA has an overall catalytic efficiency ( $k_{\text{cat}}/K_m$ ) which is  $\sim 70\%$  lower than that of the Hb<sup>+</sup> NA, indicating increased efficiency when the Hb site is present, as has been previously reported for experiments carried out with N2 modified to have Hb binding properties (5).

**Virological significance.** It has been noted in the past that the protein motifs that confer Hb activity are present in a wide range of virus subtypes which are predominantly from avian sources (4, 22). It is yet unclear whether the specific kinetic properties

**TABLE 1** Steady-state kinetic properties of N9 NAs

Substrate	NA protein	Mean $\pm$ SE $K_m$ ( $\mu\text{M}$ )	Mean $\pm$ SE $k_{\text{cat}}$ ( $\text{s}^{-1}$ )	$k_{\text{cat}}/K_m$ ( $\mu\text{M}^{-1} \text{s}^{-1}$ )
MUNANA	Hb <sup>+</sup>	$137.9 \pm 12.8$	$90.1 \pm 4.7$	0.654
MUNANA	Hb <sup>-</sup>	$134.2 \pm 12.9$	$88.6 \pm 4.7$	0.660
6SLN	Hb <sup>+</sup>	$4,179 \pm 615$	$2.2 \pm 0.2$	0.000537
3SLN	Hb <sup>+</sup>	$810 \pm 100$	$159.5 \pm 8.5$	0.197
6SLN	X-31 <sup>a</sup>	$8,070 \pm 615$	$18.5 \pm 0.9$	0.00229
3SLN	X-31 <sup>a</sup>	$562.3 \pm 20$	$97.5 \pm 1.5$	0.173
Fetuin	Hb <sup>+</sup>	$197.5 \pm 37.3$	$71.6 \pm 8.3$	0.363
Fetuin	Hb <sup>-</sup>	$346.4 \pm 95.6$	$73.0 \pm 14.5$	0.211

<sup>a</sup>The values were determined in a previous study (16).

of the N9 NA relating to the sialidase site, which favor the release rather than the cleavage of the receptor in an  $\alpha$ 2,6-linked-specific manner, are also an inherent characteristic of avian influenza viruses in general and whether these are associated with an avian host tropism.

There are potentially interesting consequences for the H7N9 viruses having Hb properties associated with the NA. The increase in overall sialic acid binding of these viruses and the enhanced binding to  $\alpha$ 2,6-linked sialic acid mediated by the sialidase site are likely advantages for the initial receptor binding to initiate infection, as the enhanced binding would increase the residence time of viruses at the surface of the cells of the human upper respiratory tract, which are rich in  $\alpha$ 2,6-linked receptors, thus likely increasing the probability of infection. This is the first reported example of the virus receptor binding specificity being determined, in part, by the NA. The N9 NA is, however, poor at cleaving  $\alpha$ 2,6-linked receptors. There are, consequently, likely to be problems relating to the inefficient release of virus from human cells after replication. This inefficient release could be a factor limiting the efficiency of aerosol transmission in ferrets (11, 23–25) and human-to-human transmission of the virus (2).

## MATERIALS AND METHODS

**Viruses.** All viruses used were constructed by reverse genetics based on a previously published system, with cDNAs for each gene segment being cloned into the pHW2000 vector (26). All viruses were generated in a 6 + 2 reassortment using the HA and NA genes from the desired virus and the remaining 6 gene segments from A/Puerto Rico/8/34(H1N1) (PR8). The H7N9 viruses were constructed using the HA and NA genes of A/Anhui/1/2013(H7N9) (Anhui13), the H1N9 viruses had the HA from PR8 and the NA from Anhui13, and the H3N9 viruses had the HA from a cell culture-propagated cultivar of A/Victoria/361/2012(H3N2) (Vic361) and the NA from Anhui13. To make viruses lacking binding via the Hb site (Hb<sup>−</sup> viruses), the NA substitution S367N was introduced into Anhui13 NA by QuikChange mutagenesis (Agilent). All viruses were rescued by the cotransfection of the 8 desired plasmids in 293T cells. H7N9 viruses were propagated in MDCK cells and H3N9 viruses were propagated in MDCK-SIAT cells in serum-free medium in the presence of 2  $\mu$ g/ml tosylsulfonyl phenylalanyl chloromethyl ketone (TPCK)-trypsin (Sigma). H1N9 viruses were propagated in 11- to 12-day-old embryonated hens' eggs. Cell culture-propagated viruses were concentrated by pelleting, and egg-propagated viruses were purified through sucrose gradients, as described previously (16). The concentration of purified/concentrated virus was determined by a solid-phase NP enzyme-linked immunosorbent assay as previously described (18). The generation, propagation, and subsequent experiments involving H7N9 and H3N9 viruses were carried out under an appropriate high biological containment level. All viruses were sequenced to ensure that no changes in the HA or NA sequence had occurred upon propagation. Plaque assays were performed with either confluent MDCK or MDCK-SIAT cells using 1.2% (wt/vol) Avicel microcrystalline cellulose and 2  $\mu$ g/ml TPCK-trypsin in 96-well plates.

**Protein expression and purification.** An expression construct containing genes encoding the Anhui13 NA ectodomain (residues 75 to 465) with an N-terminal purification tag was synthesized (GeneArt). The purification tag consisted of a hexa-His tag, a human vasodilator-stimulated phosphoprotein tetramerization domain (27, 28), and a tobacco etch virus protease cleavage site. The synthesized construct was cloned into the pFB-LIC-Bse vector with an In-Fusion cloning kit (Clontech). The substitution S367N was introduced by QuikChange mutagenesis (Agilent). Recombinant baculovirus was generated using a Bac-to-Bac system according to the manufacturer's instructions (Life Technologies). Following virus amplification, large-scale protein expression was carried out with 2.5 liters of Sf9 cells. Cells were removed by centrifugation at 72 h after infection, and the protein in the supernatant was concentrated and loaded onto a HisTALON column (Clontech). Fractions containing NA were pooled and dialyzed against 25 mM Tris-HCl, pH 8.0, 150 mM NaCl. The NA was further purified by gel filtration using a Superdex 200-pg 16/60 column (GE) in 25 mM Tris-HCl, pH 8.0, 150 mM NaCl. The gel-filtered protein was concentrated and stored in 25 mM Tris-HCl, pH 8.0, 150 mM NaCl, 4 mM CaCl<sub>2</sub>, 0.01% NaN<sub>3</sub>.

**Red blood cell capture assay.** Expressed N9 proteins were attached to His tag isolation and pulldown Dynabeads (Life Technologies). One hundred microliters of extensively washed beads was added to 50  $\mu$ l of 10  $\mu$ M N9 NA (monomeric concentration), the mixture was incubated for 30 min, and then the beads were washed to remove unbound NA. Three microliters of the beads was added to 100  $\mu$ l of 0.5% (vol/vol) turkey red blood cells (TRBCs), and the mixture was incubated at room temperature for 15 min. Samples were thoroughly mixed, and the beads were rapidly pelleted using a magnet. The supernatant was removed and allowed to settle in a microtiter plate for visual quantification, or TRBCs were lysed with SDS (final concentration, 0.5% [wt/vol]), and the released hemoglobin levels were assayed by measuring the absorbance at 540 nm.

**Virus binding studies.** Virus binding was measured as previously described (16). Briefly, binding was measured by biolayer interferometry (BLI) using an Octet Red system (Pall ForteBio Corp., Menlo Park, CA, USA). Streptavidin-coated sensors (Pall ForteBio Corp.) were loaded with biotinylated sialoglycopolymers, which consisted of 20 mol% sugar, either  $\alpha$ 2,6-sialyl-N-acetylglucosamine (6SLN) or  $\alpha$ 2,3-sialyl-N-acetylglucosamine (3SLN), attached to a polyacrylamide backbone with 5 mol% biotin (Lectinity Holdings, Moscow, Russia). Sugars were loaded at a range of concentrations for equilibrium binding assays. HA and



NA balance measurements (16) were made using sensors saturated with receptor analogues (loading  $\approx$  0.6 nm). All experiments were carried out in 10 mM HEPES, pH 7.4, 150 mM NaCl, 0.005% (vol/vol) Tween 20, 4 mM  $\text{CaCl}_2$ . When present, the NA inhibitors oseltamivir carboxylate (Roche Products Ltd., Welwyn Garden City, UK) and zanamivir (GlaxoSmithKline, Stevenage, UK) were added at concentrations of 100  $\mu\text{M}$  to inhibit NA sialidase activity. Unless stated otherwise, the virus concentration was 100 pM in all binding assays. For equilibrium binding, the total amplitude of virus binding was measured and plotted as a function of sugar loading.

**NA kinetics.** Kinetic parameters for cleavage of 2'-(4-methylumbelliferyl)- $\alpha$ -D-N-acetylneuraminic acid (MUNANA), 6SLN, and 3SLN were determined as previously described (16). Cleavage of fetuin (Sigma) was measured by using a system containing N-acetylneuraminic acid aldolase (NANA aldolase) and lactate dehydrogenase (LDH) as a reporter of released sialic acid. The 100- $\mu\text{l}$  reaction mixtures contained 0.5 U NANA aldolase (Sigma), 5 U LDH from bovine heart (Sigma), 200  $\mu\text{M}$  NADH (Sigma), and fetuin from bovine serum (Sigma) at concentrations ranging from 20 to 200  $\mu\text{M}$ . Measurements were made in 10 mM HEPES, pH 7.4, 150 mM NaCl, 0.005% (vol/vol) Tween 20, 4 mM  $\text{CaCl}_2$ . Initial reaction rates were determined by measuring the reduction in absorbance at 340 nm in a Jasco V-550 spectrophotometer using an ultramicrovolume 10-mm-path cuvette at 37°C. This change in absorbance was converted to the change in the molar concentration of NADH using its extinction coefficient ( $\epsilon = 6,220 \text{ M}^{-1} \text{ cm}^{-1}$ ).

## ACKNOWLEDGMENTS

We thank Munir Iqbal, The Pirbright Institute, UK, for the kind gift of reverse genetics plasmids for the H7 and N9 gene segments and Lauren Parker for the H3 reverse genetics plasmid. We also thank the large-scale laboratory for assistance with protein expression and purification and Evangelos Christodoulou for help developing the TRBC capture assay.

This work was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001030), the UK Medical Research Council (FC001030), and the Wellcome Trust (FC001030).

## REFERENCES

- World Health Organization. 2017. Antigenic and genetic characteristics of zoonotic influenza viruses and development of candidate vaccine viruses for pandemic preparedness. World Health Organization, Geneva, Switzerland.
- World Health Organization. 2015. WHO risk assessment of human infections with avian influenza A(H7N9) virus. World Health Organization, Geneva, Switzerland.
- Wagner R, Matrosovich M, Klenk H-D. 2002. Functional balance between haemagglutinin and neuraminidase in influenza virus infections. *Rev Med Virol* 12:159–166. <https://doi.org/10.1002/rmv.352>.
- Varghese JN, Colman PM, van Donkelaar A, Blick TJ, Sahasrabudhe A, McKimm-Breschkin JL. 1997. Structural evidence for a second sialic acid binding site in avian influenza virus neuraminidases. *Proc Natl Acad Sci U S A* 94:11808–11812. <https://doi.org/10.1073/pnas.94.22.11808>.
- Uhlendorff J, Matrosovich T, Klenk H-D, Matrosovich M. 2009. Functional significance of the hemadsorption activity of influenza virus neuraminidase and its alteration in pandemic viruses. *Arch Virol* 154:945–957. <https://doi.org/10.1007/s00705-009-0393-x>.
- Lin YP, Gregory V, Collins P, Kloess J, Wharton S, Cattle N, Lackenby A, Daniels R, Hay A. 2010. Neuraminidase receptor binding variants of human influenza A(H3N2) viruses resulting from substitution of aspartic acid 151 in the catalytic site: a role in virus attachment? *J Virol* 84:6769–6781. <https://doi.org/10.1128/JVI.00458-10>.
- Zhu X, McBride R, Nycholat CM, Yu W, Paulson JC, Wilson IA. 2012. Influenza virus neuraminidases with reduced enzymatic activity that avidly bind sialic acid receptors. *J Virol* 86:13371–13383. <https://doi.org/10.1128/JVI.01426-12>.
- Xiong X, Martin SR, Haire LF, Wharton SA, Daniels RS, Bennett MS, McCauley JW, Collins PJ, Walker PA, Skehel JJ, Gamblin SJ. 2013. Receptor binding by an H7N9 influenza virus from humans. *Nature* 499:496–499. <https://doi.org/10.1038/nature12372>.
- Yang H, Carney PJ, Chang JC, Villanueva JM, Stevens J. 2013. Structural analysis of the hemagglutinin from the recent 2013 H7N9 influenza virus. *J Virol* 87:12433–12446. <https://doi.org/10.1128/JVI.01854-13>.
- Shi Y, Zhang W, Wang F, Qi J, Wu Y, Song H, Gao F, Bi Y, Zhang Y, Fan Z, Qin C, Sun H, Liu J, Haywood J, Liu W, Gong W, Wang D, Shu Y, Wang Y, Yan J, Gao GF. 2013. Structures and receptor binding of hemagglutinins from human-infecting H7N9 influenza viruses. *Science* 342:243–247. <https://doi.org/10.1126/science.1242917>.
- Belser JA, Gustin KM, Pearce MB, Maines TR, Zeng H, Pappas C, Sun X, Carney PJ, Villanueva JM, Stevens J, Katz JM, Tumpey TM. 2013. Pathogenesis and transmission of avian influenza A (H7N9) virus in ferrets and mice. *Nature* 501:556–559. <https://doi.org/10.1038/nature12391>.
- Zhou J, Wang D, Gao R, Zhao B, Song J, Qi X, Zhang Y, Shi Y, Yang L, Zhu W, Bai T, Qin K, Lan Y, Zou S, Guo J, Dong J, Dong L, Zhang Y, Wei H, Li X, Lu J, Liu L, Zhao X, Li X, Huang W, Wen L, Bo H, Xin L, Chen Y, Xu C, Pei Y, Yang Y, Zhang X, Wang S, Feng Z, Han J, Yang W, Gao GF, Wu G, Li D, Wang Y, Shu Y. 2013. Biological features of novel avian influenza A (H7N9) virus. *Nature* 499:500–503. <https://doi.org/10.1038/nature12379>.
- Ramos I, Krammer F, Hai R, Aguilera D, Bernal-Rubio D, Steel J, García-Sastre A, Fernandez-Sesma A. 2013. H7N9 influenza viruses interact preferentially with  $\alpha$ 2,3-linked sialic acids and bind weakly to  $\alpha$ 2,6-linked sialic acids. *J Gen Virol* 94:2417–2423. <https://doi.org/10.1099/vir.0.056184-0>.
- Dortmans JCFM, Dekkers J, Wickramasinghe INA, Verheije MH, Rottier PJM, van Kuppeveld FJM, de Vries E, de Haan CAM. 2013. Adaptation of novel H7N9 influenza A virus to human receptors. *Sci Rep* 3:3058. <https://doi.org/10.1038/srep03058>.
- Xu L, Bao L, Deng W, Zhu H, Li F, Chen T, Lv Q, Yuan J, Xu Y, Li Y, Yao Y, Gu S, Yu P, Chen H, Qin C. 2014. Rapid adaptation of avian H7N9 virus in pigs. *Virology* 452-453:231–236. <https://doi.org/10.1016/j.virol.2014.01.016>.
- Benton DJ, Martin SR, Wharton SA, McCauley JW. 2015. Biophysical measurement of the balance of influenza A hemagglutinin and neuraminidase activities. *J Biol Chem* 290:6516–6521. <https://doi.org/10.1074/jbc.M114.622308>.
- Parker L, Wharton SA, Martin SR, Cross K, Lin Y, Liu Y, Feizi T, Daniels RS, McCauley JW. 2016. Effects of egg-adaptation on receptor-binding and antigenic properties of recent influenza A (H3N2) vaccine viruses. *J Gen Virol* 97:1333–1344. <https://doi.org/10.1099/jgv.0.000457>.
- Lin YP, Xiong X, Wharton SA, Martin SR, Coombs PJ, Vachieri SG, Christodoulou E, Walker PA, Liu J, Skehel JJ, Gamblin SJ, Hay AJ, Daniels RS, McCauley JW. 2012. Evolution of the receptor binding properties of the influenza A(H3N2) hemagglutinin. *Proc Natl Acad Sci U S A* 109:21474–21479. <https://doi.org/10.1073/pnas.1218841110>.
- Oh DY, Barr IG, Mosse JA, Laurie KL. 2008. MDCK-SIAT1 cells show

- improved isolation rates for recent human influenza viruses compared to conventional MDCK cells. *J Clin Microbiol* 46:2189–2194. <https://doi.org/10.1128/JCM.00398-08>.
20. Matrosovich M, Matrosovich T, Carr J, Roberts NA, Klenk H-D. 2003. Overexpression of the alpha-2,6-sialyltransferase in MDCK cells increases influenza virus sensitivity to neuraminidase inhibitors. *J Virol* 77: 8418–8425. <https://doi.org/10.1128/JVI.77.15.8418-8425.2003>.
  21. Sung JC, Van Wynsberghe AW, Amaro RE, Li WW, McCammon JA. 2010. Role of secondary sialic acid binding sites in influenza N1 neuraminidase. *J Am Chem Soc* 132:2883–2885. <https://doi.org/10.1021/ja9073672>.
  22. Kobasa D, Rodgers ME, Wells K, Kawaoka Y. 1997. Neuraminidase hemadsorption activity, conserved in avian influenza A viruses, does not influence viral replication in ducks. *J Virol* 71:6706–6713.
  23. Zhu H, Wang D, Kelvin DJ, Li L, Zheng Z, Yoon S-W, Wong S-S, Farooqui A, Wang J, Banner D, Chen R, Zheng R, Zhou J, Zhang Y, Hong W, Dong W, Cai Q, Roehrl MHA, Huang SSH, Kelvin AA, Yao T, Zhou B, Chen X, Leung GM, Poon LLM, Webster RG, Webby RJ, Peiris JSM, Guan Y, Shu Y. 2013. Infectivity, transmission, and pathology of human-isolated H7N9 influenza virus in ferrets and pigs. *Science* 341:183–186. <https://doi.org/10.1126/science.1239844>.
  24. Richard M, Schrauwen EJA, de Graaf M, Bestebroer TM, Spronken MJJ, van Boheemen S, de Meulder D, Lexmond P, Linster M, Herfst S, Smith DJ, van den Brand JM, Burke DF, Kuiken T, Rimmelzwaan GF, Osterhaus ADME, Fouchier RAM. 2013. Limited airborne transmission of H7N9 influenza A virus between ferrets. *Nature* 501:560–563. <https://doi.org/10.1038/nature12476>.
  25. Xu L, Bao L, Deng W, Dong L, Zhu H, Chen T, Lv Q, Li F, Yuan J, Xiang Z, Gao K, Xu Y, Huang L, Li Y, Liu J, Yao Y, Yu P, Li X, Huang W, Zhao X, Lan Y, Guo J, Yong W, Wei Q, Chen H, Zhang L, Qin C. 2014. Novel avian-origin human influenza A(H7N9) can be transmitted between ferrets via respiratory droplets. *J Infect Dis* 209:551–556. <https://doi.org/10.1093/infdis/jit474>.
  26. Hoffmann E, Neumann G, Kawaoka Y, Hobom G, Webster RG. 2000. A DNA transfection system for generation of influenza A virus from eight plasmids. *Proc Natl Acad Sci U S A* 97:6108–6113. <https://doi.org/10.1073/pnas.100133697>.
  27. Kühnel K, Jarchau T, Wolf E, Schlichting I, Walter U, Wittinghofer A, Strelkov SV. 2004. The VASP tetramerization domain is a right-handed coiled coil based on a 15-residue repeat. *Proc Natl Acad Sci U S A* 101:17027–17032. <https://doi.org/10.1073/pnas.0403069101>.
  28. Xu X, Zhu X, Dwek RA, Stevens J, Wilson IA. 2008. Structural characterization of the 1918 influenza virus H1N1 neuraminidase. *J Virol* 82: 10493–10501. <https://doi.org/10.1128/JVI.00959-08>.