


# Cell Host & Microbe

## Molecular Determinants for Recognition of Divergent SAMHD1 Proteins by the Lentiviral Accessory Protein Vpx

### Graphical Abstract



### Authors

David Schwefel,  
Virginie C. Boucherit, ...,  
Kate N. Bishop, Ian A. Taylor

### Correspondence

kbishop@nimr.mrc.ac.uk (K.N.B.),  
itaylor@nimr.mrc.ac.uk (I.A.T.)

### In Brief

Vpx of HIV-2 and SIV target the restriction factor SAMHD1 for degradation using the host ubiquitin ligase substrate receptor DCAF1. Schwefel et al. present the ternary complex of Vpx<sub>mnd-2</sub> with DCAF1 and the SAMHD1 N terminus and reveal how Vpx subverts host cell defenses by employing differing strategies to recruit SAMHD1 for degradation.

### Highlights

- The crystal structure of SIV<sub>mnd-2</sub> Vpx complexed with SAMHD1 and DCAF1
- Comparisons to SIV<sub>smm</sub> Vpx reveal determinants for N- versus C-terminal SAMHD1 binding
- Vpx and Vpr use conserved determinants to bind to DCAF1

### Accession Numbers

5aja



# Molecular Determinants for Recognition of Divergent SAMHD1 Proteins by the Lentiviral Accessory Protein Vpx

David Schwefel,<sup>1</sup> Virginie C. Boucherit,<sup>2</sup> Evangelos Christodoulou,<sup>1</sup> Philip A. Walker,<sup>1</sup> Jonathan P. Stoye,<sup>2,3</sup> Kate N. Bishop,<sup>2,\*</sup> and Ian A. Taylor<sup>1,\*</sup>

<sup>1</sup>Division of Molecular Structure

<sup>2</sup>Division of Virology

MRC National Institute for Medical Research, The Ridgeway, Mill Hill, London NW7 1AA, UK

<sup>3</sup>Faculty of Medicine, Imperial College London, London SW7 2AZ, UK

\*Correspondence: [kbishop@nimr.mrc.ac.uk](mailto:kbishop@nimr.mrc.ac.uk) (K.N.B.), [itaylor@nimr.mrc.ac.uk](mailto:itaylor@nimr.mrc.ac.uk) (I.A.T.)

<http://dx.doi.org/10.1016/j.chom.2015.03.004>

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## SUMMARY


The SAMHD1 triphosphohydrolase inhibits HIV-1 infection of myeloid and resting T cells by depleting dNTPs. To overcome SAMHD1, HIV-2 and some SIVs encode either of two lineages of the accessory protein Vpx that bind the SAMHD1 N or C terminus and redirect the host cullin-4 ubiquitin ligase to target SAMHD1 for proteasomal degradation. We present the ternary complex of Vpx from SIV that infects mandrills (SIV<sub>mand-2</sub>) with the cullin-4 substrate receptor, DCAF1, and N-terminal and SAM domains from mandrill SAMHD1. The structure reveals details of Vpx lineage-specific targeting of SAMHD1 N-terminal “degron” sequences. Comparison with Vpx from SIV that infects sooty mangabeys (SIV<sub>smm</sub>) complexed with SAMHD1-DCAF1 identifies molecular determinants directing Vpx lineages to N- or C-terminal SAMHD1 sequences. Inspection of the Vpx-DCAF1 interface also reveals conservation of Vpx with the evolutionally related HIV-1/SIV accessory protein Vpr. These data suggest a unified model for how Vpx and Vpr exploit DCAF1 to promote viral replication.

## INTRODUCTION

Host cell restriction factors are the first line of cellular defense against retrovirus infection. Successful viral evasion of these adaptive hurdles is essential for productive infection of new hosts and has occurred during multiple cross-species transmission events such as those that led to the emergence of pandemic human immunodeficiency virus-1 (HIV-1) (Hatzioannou and Bieniasz, 2011; Hatzioannou et al., 2014; Sharp and Hahn, 2011). Restriction factors are often interferon induced and inhibit distinct stages of the viral replication cycle (Towers and Noursadeghi, 2014; Wolf and Goff, 2008). Important examples are the TRIM5 $\alpha$ , APOBEC3 family, and Tetherin proteins, which interfere

with retroviral uncoating, reverse transcription, and budding processes, respectively (Neil et al., 2008; Sheehy et al., 2002; Stremlau et al., 2004). The sterile alpha motif domain and histidine-aspartate domain-containing protein 1 (SAMHD1) was first identified as a disease gene associated with the rare infantile encephalopathy Aicardi-Goutières syndrome (Rice et al., 2009). More recently, SAMHD1 was shown to be a factor that restricts HIV-1 replication in non-dividing myeloid-lineage cells and resting T cells (Baldauf et al., 2012; Berger et al., 2011; Descours et al., 2012; Hrecka et al., 2011; Laguetta et al., 2011). SAMHD1 is a dGTP/GTP-activated deoxynucleotide (dNTP) triphosphohydrolase (Amie et al., 2013; Goldstone et al., 2011; Ji et al., 2013; Miazzi et al., 2014; Powell et al., 2011; Zhu et al., 2013) involved in balancing cellular dNTP pools (Franzolin et al., 2013) and regulated by Cyclin A2/CDK1-dependent phosphorylation (Cribier et al., 2013; Kretschmer et al., 2015; Pauls et al., 2014; White et al., 2013). It is proposed that SAMHD1 restriction results from this triphosphohydrolase activity by reducing the cellular dNTP concentration to a level insufficient for the viral reverse transcriptase to function (Kim et al., 2012; Lahouassa et al., 2012; Rehwinkel et al., 2013; St Gelais et al., 2012). An alternative mechanism of SAMHD1 HIV-1 restriction requiring a putative nuclease activity as also been reported, but the nature of the polynucleotide substrate is disputed (Beloglazova et al., 2013; Ryoo et al., 2014; Tüngler et al., 2013).

A hallmark of most retrovirus restriction factors is the existence of viral antagonists in the form of accessory proteins (Malim and Bieniasz, 2012; Strebel, 2013). A common mechanism of action of accessory proteins is the subversion of host cell protein degradation pathways (Strebel, 2013). In particular, the host cell's Cullin-RING-type E3 ubiquitin ligases are often engaged by viral accessory proteins to induce restriction factor poly-ubiquitylation to direct proteasomal degradation (Barry and Früh, 2006). Cullin-RING ubiquitin ligases consist of a central Cullin scaffold protein, a catalytic RING subunit, and varying substrate receptors (Zimmerman et al., 2010). Owing to their modular architecture, Cullin-RING ligases allow for specific placement of a large number of substrates in the ubiquitylation zone of the catalytic subunit for efficient poly-ubiquitylation (Fischer et al., 2011; Zimmerman et al., 2010). Simian immunodeficiency virus (SIV) and HIV accessory proteins exploit these



**Figure 1. The Mandrill SAMHD1 Degron Comprises the N-Terminal and SAM Domains**

(A) Schematic of the human and mandrill SAMHD1 and the tandem NLS-EGFP degron fusion protein construct employed in fluorescent degron assays. The C- and N-terminal degron regions in SAMHD1 are highlighted in green and magenta, respectively.

(B) Stable cell lines expressing either the mandrill (mnd) SAMHD1 1–114 degron (left) or human (hs) SAMHD1 600–626 degron (right) were transduced with increasing titers of particles carrying either SIV<sub>smm</sub> Vpx (orange) or SIV<sub>mnd-2</sub> Vpx (blue) together with YFP. The level of degron (EGFP) and Vpx (YFP) expression was measured by flow cytometry. The upper panels are example FACS plots illustrating the populations of cells observed: untransduced cells (bottom right quadrant), cells transduced with a Vpx protein that can (top left quadrant) or cannot (top right quadrant) induce degradation of the degron reporter construct. The lower panels are quantification of degron reporter expression with increasing Vpx expression, demonstrating differential Vpx recognition of degron sequences. (C) Quantification of degron reporter expression in stable cells lines expressing constructs containing the indicated N-terminal regions of SAMHD1<sub>mnd</sub> after transduction with increasing amounts of SIV<sub>mnd-2</sub> Vpx; see also Figure S1.

characteristics by modifying the specificity of Cullin-RING substrate receptors (e.g., HIV-1 Vif and Vpu redirect cullin-1 and cullin-5 receptor specificity to induce APOBEC3 and Tetherin restriction factor downregulation, respectively) (Guo et al., 2014; Mitchell et al., 2009). Vpr and Vpx from HIV and SIVs target the cullin-4 ligase substrate receptor DDB1- and CUL4-associated factor 1 (DCAF1, also known as VprBP) from their host (Bergamaschi et al., 2009; Hrecka et al., 2007; Le Rouzic et al., 2007; Srivastava et al., 2008).

Restriction factors and accessory proteins are engaged in an evolutionary “molecular arms race” consisting of multiple rounds of host adaptation, virus counteraction, and host re-adaptation, resulting in accumulation of amino acid changes in restriction factor-accessory protein interaction interfaces (Daugherty and Malik, 2012). In the lentiviral Vpx/Vpr accessory proteins, analyses of positively selected residues and subsequent functional studies have demonstrated the occurrence of several significant specificity changes during adaptation to primate hosts (Laguette et al., 2012; Lim et al., 2012). It is proposed that originally a subset of Vpr proteins acquired the capability to induce cullin-4/DCAF1-dependent proteasomal degradation of SAMHD1. These SAMHD1-degrading Vpr proteins are found in the SIV<sub>syk</sub> (SIV that infects Sykes’ monkey), SIV<sub>deb</sub> (De Brazza’s monkey), and SIV<sub>agm</sub> (African green monkey) lineages. Later in its evolutionary history, a Vpr gene duplication or recombination event gave rise to Vpx, which completely took over the anti-SAMHD1 functionality and produced viruses that contain both Vpx and Vpr genes (Lim et al., 2012). In these adaptation events, the mode of SAMHD1 binding seems to have toggled between recognition sequences at the SAMHD1 N terminus and others located at the C terminus. Vpx from HIV-2, SIV<sub>smm</sub> (sooty mangabey) and SIV<sub>mac</sub> (macaque) all target the SAMHD1 C terminus,


whereas SIV<sub>mnd-2</sub> (mandrill) and SIV<sub>rcm</sub> (red-capped mangabey) Vpx induce SAMHD1 degradation dependent on the N-terminal 110 residues. The SAMHD1-degrading Vpr proteins from SIV<sub>syk</sub>, SIV<sub>deb</sub>, and SIV<sub>agm</sub> have varying specificities for the SAMHD1 N or C terminus or even target both (Fregoso et al., 2013; Spragg and Emerman, 2013).

Here, we report the crystal structure of the ternary complex of SIV<sub>mnd-2</sub> Vpx bound to the N-terminal region and SAM domain of mandrill SAMHD1 (SAMHD1<sub>mnd</sub>-NtD) and the WD40 domain of the cullin-4 substrate receptor DCAF1 (DCAF1-CtD). This structure reveals unanticipated complexity in Vpx-mediated targeting of SAMHD1. Comparison to the recently determined structure of a SIV<sub>smm</sub> Vpx/SAMHD1-CtD/DCAF1-CtD (Schwefel et al., 2014) complex provides the molecular details of the structural determinants for N-terminal versus C-terminal SAMHD1 specificity and a plausible scenario for the events occurring during evolutionary change.

## RESULTS

### Identifying the Mandrill SAMHD1 Degron

To locate the Vpx-DCAF1 recognition sequence within the SAMHD1<sub>mnd</sub> N terminus, a cell-based EGFP-degron reporter degradation assay was employed (Schwefel et al., 2014). Degron fusion proteins (Figure 1A) were constructed comprising two copies of EGFP with a nuclear localization signal (NLS) fused to either the N-terminal 114 residues of SAMHD1<sub>mnd</sub> (SAMHD1<sub>mnd</sub>-NtD) or a control C-terminal region of human SAMHD1, residues 600–626, (SAMHD1<sub>hs</sub>-CtD), which is targeted for degradation by Vpx from SIV<sub>smm</sub> (Schwefel et al., 2014). Stable cell lines expressing each reporter construct were produced and expression levels of degron fusion proteins



**Figure 2. Structure of the SIV<sub>mnd-2</sub> Vpx/DCAF1-CtD/SAMHD1<sub>mnd</sub>-NtD Complex**

(A) Schematic of components of the protein complex.

(B) Analytical gel filtration of the in vitro-assembled ternary protein complex consisting of SIV<sub>mnd-2</sub> Vpx, DCAF1-CtD and SAMHD1<sub>mnd</sub>-NtD (Peak-1). The inset shows SDS-PAGE analysis of the indicated peak fractions.

(C) Structure of the ternary complex. The backbone for each protein is shown in cartoon representation, SIV<sub>mnd-2</sub> Vpx (blue), DCAF1-CtD (gray), and SAMHD1<sub>mnd</sub>-NtD (magenta). See also Figure S2. The solvent-accessible surface is also shown for DCAF1-CtD, and a zinc ion co-ordinated by Vpx is displayed as a gray sphere, see also Figure S3. For SIV<sub>mnd-2</sub> Vpx and SAMHD1<sub>mnd</sub>-NtD, residue numbers at chain termini are indicated and secondary structure elements labeled.

were confirmed by western blot (Figure S1). Cells were then transduced with increasing amounts of a bicistronic IRES vector expressing Vpx from SIV<sub>mnd-2</sub> or SIV<sub>smm</sub> and YFP, and 48 hr later, the level of degron reporter (EGFP) and Vpx (YFP) expression was measured by flow cytometry. These data show that SIV<sub>mnd-2</sub> Vpx induces degradation of the SAMHD1<sub>mnd</sub>-NtD reporter construct but not EGFP-SAMHD1<sub>hs</sub>-CtD (Figure 1B, left panel). In contrast, SIV<sub>smm</sub> Vpx induces degradation of the SAMHD1<sub>hs</sub>-CtD reporter but not SAMHD1<sub>mnd</sub>-NtD (Figure 1B, right panel). To further delineate the SIV<sub>mnd-2</sub> SAMHD1 binding determinants, reporter constructs containing residues 1–37 and 37–114 of SAMHD1<sub>mnd</sub> were prepared. Residues 1–37 comprise an N-terminal disordered sequence containing a NLS and 37–114 constitutes the SAM domain (cp. PDB: 2E8O). How-

ever, neither fragment alone was sufficient to induce degradation of the reporter construct (Figure 1C), demonstrating that both the N-terminal NLS region and SAM domain are required for SIV<sub>mnd-2</sub> Vpx/DCAF1-mediated degradation.

#### Structure of the SIV<sub>mnd-2</sub> Vpx/SAMHD1<sub>mnd</sub>-NtD/DCAF1-CtD Ternary Complex

To gain further insights into molecular recognition in the SIV<sub>mnd-2</sub> Vpx system, SIV<sub>mnd-2</sub> Vpx, SAMHD1<sub>mnd</sub>-NtD, and DCAF1-CtD (Figure 2A) were assembled in vitro using recombinant proteins and the ternary complex purified by size-exclusion chromatography (Figure 2B). The protein complex crystallized in space group P6<sub>5</sub>22, the crystals diffracted to 2.8 Å resolution, and the structure was solved by molecular replacement using the



**Table 1. X-Ray Data Collection and Refinement Statistics**

	mnd-8498
Data collection	
Space group	P6 <sub>5</sub> 22
Cell dimensions: a, b, c (Å)	102.04, 102.04, 265.09
Cell dimensions: $\alpha$ , $\beta$ , $\gamma$ (°)	90, 90, 120
Resolution (Å)	30 (2.81) <sup>1</sup> – 2.65
$R_{\text{sym}}$ or $R_{\text{merge}}$ (%)	7.9 (145.3)
$CC_{1/2}$ <sup>2</sup>	99.9 (59.9)
$I/\sigma I$	20.70 (1.63)
Completeness (%)	99.6 (99.3)
Redundancy	9.4 (9.7)
Refinement	
Resolution (Å)	30 – 2.65
No. reflections	24,554
$R_{\text{work}}/R_{\text{free}}$	17.5/23.1
No. atoms: Protein	3,817
No. atoms: zinc ion	1
No. atoms: water	2
B factors (Å) <sup>2</sup> :	
B factors (Å) <sup>2</sup> : protein	87.6
B factors (Å) <sup>2</sup> : zinc ion	105.1
B factors (Å) <sup>2</sup> : water	71.15
rms deviations: bond lengths (Å)	0.008
rms deviations: bond angles (°)	1.183

<sup>1</sup>Values in parentheses are for highest-resolution shell. A single crystal was used for data collection.

<sup>2</sup>High-resolution cutoff based on criteria according to Karplus and Diederichs (2012).

SIV<sub>smm</sub> Vpx/SAMHD1<sub>hs</sub>-CtD/DCAF1-CtD structure (Schwefel et al., 2014) as a search model. Details of the data collection, structure determination, map quality, and refinement are presented in Table 1 and Figure S2.

DCAF1-CtD consists of a typical WD40-repeat seven-bladed  $\beta$ -propeller fold approximately 45 Å diameter and 25 Å in height (Figure 2C). SIV<sub>mnd-2</sub> Vpx comprises an N-terminal extended arm that precedes three  $\alpha$  helices. The side chains of amino acid residues H35 at the C terminus of helix  $\alpha$ 1 together with H78 and C83 at the C terminus of helix  $\alpha$ 3 constitute a zinc-binding motif that links the helices together, similar to that observed in SIV<sub>smm</sub> Vpx (Schwefel et al., 2014). However, in contrast the fourth zinc coordinating ligand in SIV<sub>mnd-2</sub> Vpx is a water molecule rather than an additional cysteine C-terminal to C83 as occurs in SIV<sub>smm</sub> Vpx (Figure S3). In the structure, two sections of the SAMHD1<sub>mnd</sub>-NtD are visible. Residues 1–22 are in an extended conformation, and residues 34–109 fold into the SAM domain, which consists of four  $\alpha$  helices packed together by interaction of inward-facing hydrophobic amino acid side chains. The ternary complex is built up by three major protein interaction interfaces. These are (i) a Vpx-DCAF1 interface on top and side of the DCAF1 propeller, (ii) a combined interface between the N-terminal extended segment of SAMHD1-NtD, DCAF1 and the Vpx on the side of DCAF1, and (iii) contacts between the SAM domain and Vpx on top of DCAF1.


### The DCAF1 Binding Mode Is Conserved in Vpx Proteins

The extensive Vpx-DCAF1 interface comprises four contact regions with a total surface area of 1,600 Å<sup>2</sup> (Figure 3; Table S1A). The first region comprises the C-terminal section of  $\alpha$ 2 together with first half of  $\alpha$ 3 of Vpx. On one face, apolar amino acid side chains from both  $\alpha$ 2 and  $\alpha$ 3 pack against a cluster of hydrophobic amino acids located in the “tall” loop that connects  $\beta$ 2 to  $\beta$ 3 in WD40 repeat 7 on the upper surface of DCAF1-CtD (Figure 3B, upper panel). On the other face, a network of hydrogen bonds and salt bridges connects Vpx residues Y62, Y65, R66, and K73 at the N terminus of  $\alpha$ 3 to acidic side chains E1191, D1092, and E1193 in the “acidic” loop of DCAF1 that intersperses WD40 repeats 7 and 1 (acidic loop, Figure 3B, lower panel). A second region of interaction involves the carboxy terminus of Vpx  $\alpha$ 3, which packs into a radial groove on the top side of DCAF1 lined by amino acids from WD40 repeats 1, 2, 4, 6, and 7. Interactions here include both the packing of hydrophobic residues as well as specific hydrogen bonding between the side-chains of Vpx Y76 and Q81 and the DCAF1 protein backbone (Figure 3C). The third binding interface involves residues at the extreme amino terminal of the Vpx N-terminal extended arm that inserts into a cavity between WD40 repeats 1 and 2 on the underside of DCAF1. Here, Vpx A2, E3 and E7 make main chain and side chain hydrogen-bonds to DCAF1 residues S1102, R1106, Y1131, and S1168 while the interposing Vpx residues A5 and P6 project into the hydrophobic cavity (Figure 3D). The remaining site of interaction involves contacts between residues in the WD40 repeat 2 of DCAF1 and residues in both Vpx  $\alpha$ 1 and  $\alpha$ 3. These include hydrophobic interactions between DCAF1 T1155 and W1156 and apolar side chains displayed on the underside of Vpx  $\alpha$ 1 and  $\alpha$ 3 and Vpx H72 on  $\alpha$ 3 that makes a hydrogen bond with mainchain carbonyl of N1135 of DCAF1 (Figure 3E).

Structural alignment of the DCAF1-CtD and Vpx molecules in the SIV<sub>mnd-2</sub> and SIV<sub>smm</sub> complexes (Schwefel et al., 2014) reveals that in both instances they adopt very similar conformations. The root-mean-square deviations (RMSDs) are 0.65 Å (305 aligned residues) for the DCAF1-CtD molecules and 1.24 Å for Vpx (80 aligned residues) (Figure S4A). Moreover, detailed comparison of SIV<sub>mnd-2</sub> and SIV<sub>smm</sub> Vpx amino acid side chains that contact DCAF1 shows that nearly all are at least type-conserved (Figures S4B–S4E). A notable exception is T47 in  $\alpha$ 2 that is exchanged to arginine in SIV<sub>smm</sub> Vpx and is concomitant with a small movement of  $\alpha$ 2 and differences in conformation of the interacting DCAF1 tall loop between the two complexes (Figure S4B, upper panel). However, residues on the upper face of  $\alpha$ 2 also interact with the SAMHD1<sub>mnd</sub> SAM domain (see below) so the cause of the  $\alpha$ 2 shift is unclear. Therefore, given that the mode of the N-terminal arm -  $\alpha$ 2 -  $\alpha$ 3 Vpx association with DCAF1 is a common feature of SIV<sub>mnd-2</sub> and SIV<sub>smm</sub> complexes and that nearly all the DCAF1-interacting residues are type-conserved in HIV-2/SIV Vpx proteins (Figure S5), it is likely that the same principles of DCAF1 binding apply to both the SIV<sub>mnd-2</sub> and SIV<sub>smm</sub> Vpx lineages.

### A Combined SIV<sub>mnd-2</sub> Vpx/SAMHD1<sub>mnd</sub>(1–22)/DCAF1-CtD Ternary Interface

In the complex, the N-terminal NLS region (residues 1–22) of SAMHD1<sub>mnd</sub> comprises an extended chain that binds to the




**Figure 3. The SIV<sub>mnd-2</sub> Vpx/DCAF1-CtD Interaction Interface**

(A) Overview of Vpx/DCAF1-CtD interactions. Molecules are in the same representation as Figure 2, the WD40  $\beta$ -propeller repeats of DCAF1-CtD are numbered. (B–E) Details of Vpx/DCAF1-CtD interactions in the regions boxed in (A). Residues that make interactions are shown in stick representation, dashed lines indicate hydrogen bonds and salt bridges, see also Figure S4 and Table S1.

edge of DCAF1 in an interface that buries 500 Å<sup>2</sup> of solvent accessible surface (Figure 4A; Table S1B). Here, residues D5, D7, and Q8 make extensive sidechain and mainchain hydrogen bonds to residues in the interspersing loops of DCAF1 WD40 repeat-1. In addition, SAMHD1 residues R12 and R14, which are part of the SAMHD1 NLS (Brandariz-Núñez et al., 2012; Hofmann et al., 2012), make salt bridges to E1091 and D1092 in the DCAF1 acidic loop. The ternary interaction is completed by a 750 Å<sup>2</sup> SAMHD1<sub>mnd</sub>-SIV<sub>mnd-2</sub> Vpx interface that includes multiple interactions with Vpx residues from the N-terminal arm,  $\alpha$ 1, and the  $\alpha$ 2- $\alpha$ 3 loop (Figure 4B; Table S1C). One feature of this interface is a “trapped” region where the Vpx N-terminal arm wraps over SAMHD1 packing it against DCAF1. Within the trapped region, the main chains of SAMHD1 and Vpx interact through hydrogen bonding between E10, G13, and E14 of Vpx

and Q2 and S10 of SAMHD1. In addition, the main chain of Vpx G13, and E14 also make interactions with the SAMHD1 D7 and S10 side chains. C-terminal to the trapped region there are further interactions where Vpx V15, L17, and W20 create a hydrophobic pocket at the N terminus of  $\alpha$ 1 that SAMHD1 P9 inserts into. Additionally, SAMHD1 NLS residue R12 makes a hydrogen bond to Vpx Y65, SAMHD1 F15 stacks with Vpx R58, and SAMHD1 R20 makes an electrostatic interaction with Vpx D54 (Figure 4B).

The functional importance of the Vpx trapping of the SAMHD1 N-terminal region was revealed in an EGFP-degron fusion protein degradation assay. Figure 4C shows that SIV<sub>mnd-2</sub> Vpx induces efficient degradation of constructs containing SAMHD1 amino acids 1–114 and 5–114. In contrast, EGFP-SAMHD1 (10–114), which lacks D7 and Q8 that make hydrogen bonds to



**Figure 4. The Combined SIV<sub>mnd-2</sub> Vpx/DCAF1-CtD/SAMHD1<sub>mnd</sub>-NtD Ternary Interface**

(A) Interface of DCAF1-CtD and SAMHD1<sub>mnd</sub> NLS region.  $\beta$  strands from WD40 repeat-1 and -2 of DCAF1-CtD are shown in gray cartoon representation. The N-terminal NLS region of SAMHD1<sub>mnd</sub> is shown in magenta. Residues making contacts at the interface are shown in stick representation with hydrogen bonding and salt bridge interactions displayed as dashed lines.

(B) The Vpx-SAMHD1<sub>mnd</sub> NLS region interface. The ternary complex is displayed (central) in the same orientation and representation as Figure 2. Details of SAMHD1-Vpx interactions within the boxed region are shown left and right. Residues in SAMHD1 (magenta) and Vpx (blue) making contacts at the interface are shown in stick representation with hydrogen bonding and salt bridge interactions displayed as dashed lines.

(C) Quantification of reporter expression in stable cell lines containing degron constructs with the indicated N-terminal truncations of SAMHD1<sub>mnd</sub>. Cells were transduced with increasing titers of particles expressing SIV<sub>mnd-2</sub> Vpx together with YFP and degron reporter expression (EGFP) measured by flow cytometry.

Vpx and DCAF1 within the trapped region, is not degraded by Vpx indicating the requirement of the trapping interaction for degron activity.


#### The SIV<sub>mnd-2</sub> Vpx/ SAMHD1<sub>mnd</sub> SAM Domain Interface

In addition to the ternary interface that “traps” the SAMHD1<sub>mnd</sub> N terminus within the complex a further DCAF1-independent interaction between the SAM domain of SAMHD1<sub>mnd</sub>-NtD and SIV<sub>mnd-2</sub> Vpx is also observed. This interface has a buried surface area of 580 Å<sup>2</sup> and comprises residues that project upward from Vpx  $\alpha$ 2 that contact SAMHD1 residues N-terminal to the SAM domain helix 1 as well as those on the underside of the SAM domain in helices 1 and 2 (Figure 5A). The center of the interface comprises a hydrophobic core created by Vpx residues P38, L41, F42, W45, and V49 and SAMHD1 residues P34, V36, L38, and F52 (Figure 5A, upper panel). In addition to this hydrophobic patch, there are electrostatic interactions between SAMHD1 R55 and Vpx D54 (Figure 5A, upper panel), SAMHD1 R69 and Vpx E50 (Figure 5A, lower panel), as well as hydrogen bonding from Vpx E39 to the backbone of the first turn of SAM domain helix 1 (Figure 5B, lower panel). The contribution of res-

idues in the SAM domain-Vpx interface to degron function was assessed by introduction of alanine mutations at SAMHD1 residues R55 and R69 in the EGFP-fusion reporter. Both residues make extensive electrostatic interactions at the interface, and the alanine substitutions result in a loss of reporter degradation (Figure 5B). R69 is also among a group of residues in the N-terminal region of SAMHD1 that have undergone positive evolutionary selection (Lim et al., 2012) indicative of an interaction hotspot between restriction factor and viral antagonist, further underlining the importance of Vpx-SAM domain contacts for induction of SAMHD1 proteasomal degradation.

Comparison of the sequences of human and mandrill SAMHD1-NtD regions reveals 13 amino-acid changes, nine in the SAM domain and four in the NLS region. Of these, only two make side chain contacts in the mandrill ternary complex, F15 in the NLS region and F52 in the SAM domain. In human SAMHD1, these residues are replaced by non-conservative cysteine and serine substitutions (Figure S6). In the mandrill complex, both phenylalanine residues make interactions, F15 stacks against the side chain of R58 in the Vpx  $\alpha$ 2- $\alpha$ 3 loop (Figure 4B, left panel), and F52 is central to the hydrophobic SAM





**Figure 5. SAM Domain-SIV<sub>mnd-2</sub> Vpx Interactions**

(A) The SAMHD1<sub>mnd</sub> SAM domain-SIV<sub>mnd-2</sub> Vpx interface. The SAMHD1<sub>mnd</sub> SAM domain (magenta) and SIV<sub>mnd-2</sub> Vpx (blue) are shown in cartoon representation. Residues making interactions are shown in stick representation with hydrogen bonding and salt bridges displayed as dashed lines.

(B and C) Quantification of reporter expression in cells stably expressing SAMHD1<sub>mnd</sub> (1-114) degon reporter constructs with (B) SAM domain point mutations and (C) mandrill to human SAMHD1 amino acid substitutions. Cells were transduced with increasing titers of particles expressing SIV<sub>mnd-2</sub> Vpx, and the level of transduction (YFP) and degon reporter (EGFP) expression was measured by flow cytometry.

domain-Vpx interaction (Figure 5A). In order to assess their role in the specificity of the Vpx-SAMHD1 interaction, F15 and F52 were mutated to cysteine and serine that are found in human

SAMHD1 and their contribution tested in the degon assay (Figure 5C). These data show the F15C mutation renders SAMHD1<sub>mnd</sub> resistant to SIV<sub>mnd-2</sub> Vpx-induced proteasomal degradation but that surprisingly the F52S mutation only slightly reduces degradation of the degon reporter protein. Nevertheless, introduction of the reverse mutations C15F and S52F in human SAMHD1 has previously been shown to render human SAMHD1 sensitive to SIV<sub>mnd-2</sub> Vpx-mediated degradation (Wei et al., 2014). Therefore, taken together, these data support the idea that amino acid changes in primate SAM domain and NLS regions determine the lineage-specific formation of a Vpx-SAMHD1 interaction interface.


### Specificity of Vpx-SAMHD1-NtD and -CtD Interactions

Despite similar overall structure and an identical mode of DCAF1 interaction (Figure S4), the SIV<sub>mnd-2</sub> Vpx/DCAF1 assembly recognizes the SAMHD1<sub>mnd</sub>-NtD, while the HIV-2/SIV<sub>mac</sub>/SIV<sub>smm</sub>-type Vpx/DCAF1 are specific for the SAMHD1 C terminus. Comparison of the SIV<sub>mnd-2</sub> system presented here with the previously determined SIV<sub>smm</sub>/DCAF1/SAMHD1-CtD assembly (Schwefel et al., 2014) reveals the structural determinants for these fundamentally different specificities. In both complexes, a Vpx N-terminal arm and residues at the N terminus of  $\alpha 1$  make contacts with either the amino-terminal region in SAMHD1<sub>mnd</sub> or the CtD region of SAMHD1<sub>hs</sub>. The location of this contact on the DCAF1 disk is similar in both combined ternary interfaces (Figure 6A). However, the SAMHD1<sub>mnd</sub> N terminus makes additional contacts and occupies a larger surface area on Vpx and DCAF1, 750 Å<sup>2</sup> and 500 Å<sup>2</sup> versus 710 Å<sup>2</sup> and 210 Å<sup>2</sup> for the SAMHD1<sub>hs</sub> C terminus. In addition, in the mandrill complex, the Vpx arm folds over the first 20 residues of SAMHD1<sub>mnd</sub> trapping it against DCAF1, while the human SAMHD1 C-terminal peptide associates rather peripherally to the DCAF1/Vpx interface (Figure 6A).

In this context, it is important to note that Vpx amino acid residues at the N terminus of  $\alpha 1$  that contribute to SAMHD1 binding differ significantly between SIV<sub>smm</sub> and SIV<sub>mnd-2</sub> type Vpx proteins constituting a variable region (VR1) (Figure S5). In the SIV<sub>smm</sub> complex, VR1 side chains of the conserved GEET-motif form salt bridges with basic residues in SAMHD1-CtD (Schwefel et al., 2014). However, in SIV<sub>mnd-2</sub> Vpx, VR1 comprises a proline (P9<sup>mnd-2</sup>) followed by two glycine residues interspersed with a small hydrophobic amino acid and then a conserved acidic and hydrophobic residue (<sup>9</sup>P[x]GAG[D/E]V/A<sup>15</sup>). This configuration specifically accommodates the SAMHD1<sub>mnd</sub>-NtD and sandwiches it between Vpx and the DCAF1 disk. The importance of this VR1 interaction for Vpx-mediated proteasomal degradation of SAMHD1<sub>mnd</sub> is demonstrated in EGFP-degon fusion protein assays where substitution of valine by bulky tryptophan at position 15 of SIV<sub>mnd-2</sub> Vpx substantially reduces degon reporter protein degradation (Figure 6B). In addition, analysis of the clustering of VR1 sequences (Figure S5) shows that SIV<sub>rcm</sub> and SIV<sub>drl-1</sub> Vpx that are SAMHD1-NtD specific also contain a SIV<sub>mnd-2</sub> type VR1, supporting the idea that VR1 is a major determinant for discrimination between SAMHD1 N-terminal and C-terminal target sequences.

As well as VR1, which confers NtD and CtD degon specificity to Vpx, a second region, VR2, in the  $\alpha 2$ - $\alpha 3$  loop and N terminus of  $\alpha 3$  is conserved among Vpx but not in Vpr (Figure S5). Here,





**Figure 6. Comparison to the SIV<sub>smm</sub> Vpx/DCAF1-CtD/SAMHD1<sub>hs</sub>-CtD Structure**

(A) Superposition of the SIV<sub>mnd-2</sub> Vpx/DCAF1-CtD/SAMHD1<sub>mnd</sub>-NtD and SIV<sub>smm</sub> Vpx/DCAF1-CtD/SAMHD1<sub>hs</sub>-CtD ternary complexes. Cartoons are colored as Figure 2 with the addition of SIV<sub>smm</sub> Vpx (orange) and SAMHD1<sub>hs</sub>-CtD (green). For clarity, only DCAF1-CtD from the mandrill complex is shown.

(B and C) Quantification of reporter expression in cells stably expressing the SAMHD1<sub>mnd</sub> (1–114) degron reporter construct after transduction with increasing titers of particles expressing SIV<sub>mnd-2</sub> Vpx with mutations in (B) the N-terminal arm (V15W) and in (C) the di-tyrosine motif (Y62A/Y65A). The level of Vpx transduction (YFP) and degron reporter (EGFP) expression was measured by flow cytometry.

(D) Detailed view of the conserved VR2 Vpx di-tyrosine motif interaction with the DCAF1-CtD acidic loop and basic motifs of the SAMHD1<sub>mnd</sub>-NtD (left) and SAMHD1<sub>hs</sub>-CtD (right) degrons; see also Figure S5. Residues that contribute to the ternary interface are shown in stick representation with hydrogen bonding and salt bridge interactions displayed as dashed lines; see also Figure S6.

a di-tyrosine motif (Y62<sup>mnd-2</sup>/Y66<sup>smm</sup> and Y65<sup>mnd-2</sup>/Y69<sup>smm</sup>) is present, which engages the DCAF1 acidic loop (Figures 3B and S4B), providing a hydrophobic platform for binding of SAMHD1 residues (P13<sup>mnd</sup> or V618<sup>hs</sup>/F621<sup>hs</sup>) and for positively charged amino acids (R12<sup>mnd</sup>/R14<sup>mnd</sup> or K622<sup>hs</sup>) to form salt bridges to the DCAF1 acidic loop (Figure 6D). The importance of the Vpx di-tyrosine pair at the center of a ternary DCAF1-degion interaction was demonstrated in the degon assay where introduction of a double tyrosine to alanine mutant into SIV<sub>mnd-2</sub> Vpx abolished reporter degradation (Figure 6C). Similarly, mutations of the basic residues that make salt bridges to the DCAF1 acidic loop also prevent degon reporter degradation (Schwefel et al., 2014) and in a pigtail macaque model SIV<sub>mne</sub>, a Vpx Y66A, Y69A, Y71A triple mutant causes a reduction of viral load (Belsham et al., 2012). Taken together, these observations suggest that the VR2 region contains common binding determinants utilized in both SIV<sub>mnd-2</sub> and SIV<sub>smm</sub> type Vpx molecules for recognition of SAMHD1-NtD and SAMHD1-CtD degons.

## DISCUSSION

The current properties of accessory proteins Vpr and Vpx likely reflect a lengthy history of conflict between lentiviruses and their hosts. Eight different lineages of exogenous primate lentivirus including HIV-1 can be identified. All encode a Vpr that induces host G2/M cell-cycle arrest (Berger et al., 2015), probably by tar-

geting a still poorly defined host cell protein for degradation through the proteasomal pathway or through interaction with proteins of the DNA damage pathway (Laguette et al., 2014). Vpr from two viral clades, typified by SIV<sub>agm</sub> and SIV<sub>syk</sub>, is also able to target SAMHD1 for degradation, facilitating infection of some types of non-dividing cells. In two further clades, both Vpr and Vpx are present with Vpr retaining only cell-cycle arrest activity and Vpx only inducing SAMHD1 degradation.

Phylogenetic reconstructions suggest that the Vprs from SIV<sub>agm</sub> and SIV<sub>syk</sub> acquired anti-SAMHD1 activity prior to the molecular events leading to the generation of Vpx (Lim et al., 2012). Moreover, it appears that once Vpr acquires anti-SAMHD1 activity, it does not lose it. This would imply that viruses containing both Vpr and Vpx arose by the recombination of two viruses; the first encoding an ancestral Vpr with the dual ability to cause cell-cycle arrest and degrade SAMHD1 and the second with a Vpr that had not acquired the capacity to induce SAMHD1 degradation. This allowed the ancestral, dual function Vpr to convert to Vpx and maintain SAMHD1 degradation activity but lose the cell-cycle arrest function provided by the second Vpr. Evaluation of our crystal structures is most informative in understanding these evolutionary changes and specifically the interaction between (i) Vpr/x and DCAF1 and (ii) between the two classes of Vpx and their SAMHD1 targets.

Amino acid residues that bind to DCAF1 and those that coordinate the bound zinc are highly conserved in all lentiviral Vpr/Vpx proteins (Figure S5), suggesting that the overall structure of the protein as well as mode of DCAF1 binding have been conserved during Vpr/Vpx evolution. This implies that the

HIV-1 Vpr/DCAF1 association is likely to be similar to the one observed in the SIV<sub>mnd-2</sub> and SIV<sub>smm</sub> (Schwefel et al., 2014) Vpx/DCAF1 crystal structures (Figure S4). Therefore, these structures might provide suitable starting points for rational design of compounds interfering with HIV-1 Vpr function even in the absence of definitive information concerning its cellular target. A notable exception is Vpr from SIV<sub>deb</sub> that induces SAMHD1 degradation apparently without binding to DCAF1 (Berger et al., 2015), and it will be interesting to determine whether it utilizes an alternative pathway or different receptor protein to target SAMHD1.

Vpx from SIV<sub>smm</sub> and SIV<sub>mnd-2</sub> have been shown to interact with the C- and N-terminal regions of SAMHD1, respectively (Fre-goso et al., 2013). Comparison of Vpx/SAMHD1/DCAF1 ternary complexes reveals that this discrimination of SAMHD1 N- and C-terminal degrons by the two classes of Vpx is determined both by polymorphisms in the SAMHD1 amino-terminal region (Figure S6) and by specific sequences in the VR1 region of Vpx (Figures 6 and S5). By contrast, in both classes of Vpx the VR2 region contains a conserved di-tyrosine pair that interacts with the DCAF1 acidic loop to provide a hydrophobic platform for N- and C-terminal SAMHD1 degrons. In this way, basic residues from both degron types are positioned for electrostatic interactions with the DCAF1 acidic loop. Interestingly, only the second tyrosine is present in SIV<sub>agm</sub> Vpr proteins, which are incapable of inducing human SAMHD1 degradation, while the position corresponding to the first tyrosine is occupied by a glutamate residue (Figure S5). Gain of the bi-tyrosine motif might therefore have been a critical step in the development of Vpx allowing multiple modes of binding SAMHD1 while at the same time resulting in loss of the Vpr capacity to cause cell-cycle arrest.

A third highly variable region (VR3) in Vpr/Vpx proteins is directly C-terminal to the fourth conserved zinc-coordinating residue (Figure S5) that in both SIV<sub>smm</sub> and SIV<sub>mnd-2</sub> type Vpx contains proline-rich stretches of variable length. In the corresponding SAMHD1/Vpx/DCAF1 crystal structures, VR3 is either disordered (this study) or involved in crystal contacts (Schwefel et al., 2014), strongly suggesting that the region is not involved in SAMHD1 degron recognition. In agreement, SIV<sub>mac</sub> Vpx lacking the poly-proline tail is still able to assemble into a ternary SAMHD1/Vpx/DCAF1 complex and to induce cullin-4-dependent SAMHD1 poly-ubiquitination (Ahn et al., 2012) and degradation (Belshan et al., 2012). Functional studies suggest involvement of VR3 in nuclear localization of Vpx and Vpr (Pancio et al., 2000; Zhou et al., 1998) with consequences for viral replication (Belshan et al., 2012), but notably, mutations in this region also influence protein expression levels significantly (Miyake et al., 2014).

Predictions regarding DCAF1/Vpr-mediated recruitment of target protein(s) that result in G2/M cell-cycle arrest remain speculative. In HIV-1, VR2 differs significantly from other Vpr and Vpx proteins, suggesting that in contrast to the Vpx di-tyrosine/DCAF1 acidic loop/SAMHD1 interaction, other principles may apply to HIV-1 Vpr target recognition. By contrast, VR1 of HIV-1 Vpr contains a <sup>12</sup>REP[F/Y/W][D/N]<sup>16</sup> motif that is also conserved in SIV<sub>agm</sub> Vpr proteins (Figure S5), suggesting similarities in VR1-mediated SIV<sub>agm</sub> and HIV-1 Vpr target recruitment. Therefore, considering recent evidence for HIV-1 Vpr/DCAF1 association with components of the SLX4/MUS81/EME1 DNA-

repair complex (Berger et al., 2015; Laguetta et al., 2014), structural analysis of SIV<sub>agm</sub> Vpr/DCAF1 together with the cognate SAMHD1<sub>agm</sub> will be of interest to further delineate the degron and/or target sequence(s) recognized by HIV-1 Vpr.

## EXPERIMENTAL PROCEDURES

### Degron Assay

Degron reporter constructs were generated by replacing the human SAMHD1-CtD degron sequence in pCMS28-NLS-EGFP-SAMHD1-CtD with N-terminal sequences from SAMHD1<sub>mnd-2</sub>. Point mutations were created by site-directed mutagenesis. Virus-like particles (VLPs) were generated by co-transfecting 293T cells with pVSVG, pKB4 and pCMS28-NLS-EGFP-SAMHD1. Stable cell lines were produced by transduction of *Mus dunni* cells followed by puromycin selection. Expression of degron constructs was assessed using western blotting with anti-EGFP antibodies.

SIV<sub>smm</sub> or SIV<sub>mnd-2</sub> Vpx sequences were transferred into pLgatewayYFP (Gateway LR clonase™ II, Invitrogen) to create bicistronic Vpx-IRES-YFP expression constructs. Point mutations were created by site-directed mutagenesis. VLPs expressing Vpx-IRES-YFP were generated by co-transfecting 293T cells with pVSV-G, pKB4, and pLgatewayYFP-Vpx.

Parental *Mus dunni* or stable cell lines expressing degron reporters were seeded at  $5 \times 10^4$  cells per well in a 24-well plate 1 day prior to infection with 2-fold serial dilutions of Vpx-YFP VLPs. After 48 hr, the percentage of EGFP- and YFP-positive cells was determined by flow cytometry.

### Protein Expression and Purification

The nucleotide sequences coding for SIV<sub>mnd-2</sub> Vpx isolate 5440 and amino acid residues 1–114 of SAMHD1<sub>mnd-2</sub> were inserted into pET-49b and pET-52b (Merck Millipore) expression plasmids respectively to generate N-terminally GST-tagged and Strep-II-tagged fusion proteins. SAMHD1 (1–114) was expressed in *E. coli* strain Rosetta 2 and purified using Strep-Tactin affinity followed by size-exclusion chromatography. SIV<sub>mnd-2</sub> Vpx was captured from the bacterial lysate onto Glutathione Sepharose (GSH-Sepharose) (GE Healthcare) prior to an assembly reaction (see below). His-tagged DCAF1, residues 1,058–1,396 (DCAF1-CtD), was expressed in insect cells and purified using Ni-NTA Sepharose and size-exclusion chromatography.

For assembly, the GST-SIV<sub>mnd-2</sub> Vpx bound beads were resuspended and incubated with 1 mg of DCAF1-CtD, an equimolar amount of SAMHD1<sub>mnd-2</sub> NtD, and 1 mg of HRV-3C protease (GE Healthcare) overnight at 4°C. After removal of the beads by centrifugation, the eluted ternary complex was further purified by size-exclusion chromatography. Details of protein expression constructs, protein production, and storage are provided in Supplemental Experimental Procedures.

### Crystallization and Structure Solution

Crystals of the SIV<sub>mnd-2</sub> Vpx/SAMHD1<sub>mnd-2</sub>NtD/DCAF1CtD complex were grown using the hanging drop vapor diffusion at 18°C in 2 µl droplets comprising 1 µl complex (6.34 mg/ml) and 1 µl of reservoir solution (0.16 M Trisodium Citrate-HCl [pH 5.2] and 4% PEG 6000). X-ray diffraction data were collected on beamline I04 at the Diamond Light Source, UK at a wavelength of 0.97965 Å. The structure was solved by molecular replacement using the previously determined DCAF1-CtD structure, and a homology model constructed with the previously determined SIV<sub>smm</sub> Vpx as template (PDB: 4CC9). Details of crystallization, data collection, processing, and structure solution are provided in Supplemental Experimental Procedures.

### Multiple Sequence Alignment

Amino acid sequences were aligned using the ClustalW server and adjusted manually. NCBI accession numbers for Vpr, Vpx, and SAMHD1 are provided in Supplemental Experimental Procedures.

### ACCESSION NUMBERS

The coordinates and structure factors of the SIV<sub>mnd-2</sub>-Vpx/SAMHD1<sub>mnd-2</sub>NtD/DCAF1-CtD ternary complex have been deposited in the Protein Data Bank (PDB: 5aja).

## SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures, three table, and Supplemental Experimental Procedures and can be found with this article online at <http://dx.doi.org/10.1016/j.chom.2015.03.004>.

## AUTHOR CONTRIBUTIONS

D.S., V.C.B., E.C., and P.A.W. performed experiments. D.S., V.C.B., E.C., P.A.W., J.P.S., K.N.B., and I.A.T. contributed to experimental design, data analysis, and manuscript writing. The authors declare no competing financial interests.

## ACKNOWLEDGMENTS

We gratefully acknowledge Diamond Light Source for synchrotron access. This work was supported by the UK Medical Research Council file references U117512710 (J.P.S.), U117592729 (K.N.B.), and U117565647 (I.A.T.); The Wellcome Trust, ref 085955 (K.N.B.); an EMBO long-term fellowship co-funded by the European Commission Marie Curie Actions (EMBOCOFUND2010, GA-2010-267146) (D.S.); and an amfAR Mathilde Krim Fellowship in Basic Biomedical Research ref 108992-57-RKHF (D.S.).

Received: January 8, 2015

Revised: January 31, 2015

Accepted: February 27, 2015

Published: April 8, 2015

## REFERENCES

Ahn, J., Hao, C., Yan, J., DeLucia, M., Mehrens, J., Wang, C., Gronenborn, A.M., and Skowronski, J. (2012). HIV/simian immunodeficiency virus (SIV) accessory virulence factor Vpx loads the host cell restriction factor SAMHD1 onto the E3 ubiquitin ligase complex CRL4DCAF1. *J. Biol. Chem.* 287, 12550–12558.

Amie, S.M., Bambara, R.A., and Kim, B. (2013). GTP is the primary activator of the anti-HIV restriction factor SAMHD1. *J. Biol. Chem.* 288, 25001–25006.

Baldauf, H.M., Pan, X., Erikson, E., Schmidt, S., Daddacha, W., Burggraf, M., Schenkova, K., Ambiel, I., Wabnitz, G., Gramberg, T., et al. (2012). SAMHD1 restricts HIV-1 infection in resting CD4(+) T cells. *Nat. Med.* 18, 1682–1687.

Barry, M., and Früh, K. (2006). Viral modulators of cullin RING ubiquitin ligases: culling the host defense. *Sci. STKE* 2006, pe21.

Beloglazova, N., Flick, R., Tchigvintsev, A., Brown, G., Popovic, A., Nocek, B., and Yakunin, A.F. (2013). Nuclease activity of the human SAMHD1 protein implicated in the Aicardi-Goutières syndrome and HIV-1 restriction. *J. Biol. Chem.* 288, 8101–8110.

Belshan, M., Kimata, J.T., Brown, C., Cheng, X., McCulley, A., Larsen, A., Thippeshappa, R., Hodara, V., Giavedoni, L., Hirsch, V., and Ratner, L. (2012). Vpx is critical for SIVmac infection of pigtail macaques. *Retrovirology* 9, 32.

Bergamaschi, A., Ayinde, D., David, A., Le Rouzic, E., Morel, M., Collin, G., Descamps, D., Damond, F., Brun-Vezinet, F., Nisole, S., et al. (2009). The human immunodeficiency virus type 2 Vpx protein usurps the CUL4A-DDB1 DCAF1 ubiquitin ligase to overcome a postentry block in macrophage infection. *J. Virol.* 83, 4854–4860.

Berger, A., Sommer, A.F., Zwarg, J., Hamdorf, M., Welzel, K., Esly, N., Panitz, S., Reuter, A., Ramos, I., Jattiani, A., et al. (2011). SAMHD1-deficient CD14+ cells from individuals with Aicardi-Goutières syndrome are highly susceptible to HIV-1 infection. *PLoS Pathog.* 7, e1002425.

Berger, G., Lawrence, M., Hué, S., and Neil, S.J. (2015). G2/M cell cycle arrest correlates with primate lentiviral Vpr interaction with the SLX4 complex. *J. Virol.* 89, 230–240.

Brandariz-Núñez, A., Valle-Casuso, J.C., White, T.E., Laguerre, N., Benkirane, M., Brojatsch, J., and Diaz-Griffero, F. (2012). Role of SAMHD1 nuclear localization in restriction of HIV-1 and SIVmac. *Retrovirology* 9, 49.

Cribier, A., Descours, B., Valadão, A.L., Laguerre, N., and Benkirane, M. (2013). Phosphorylation of SAMHD1 by cyclin A2/CDK1 regulates its restriction activity toward HIV-1. *Cell Rep.* 3, 1036–1043.

Daugherty, M.D., and Malik, H.S. (2012). Rules of engagement: molecular insights from host-virus arms races. *Annu. Rev. Genet.* 46, 677–700.

Descours, B., Cribier, A., Chable-Bessia, C., Ayinde, D., Rice, G., Crow, Y., Yatim, A., Schwartz, O., Laguerre, N., and Benkirane, M. (2012). SAMHD1 restricts HIV-1 reverse transcription in quiescent CD4(+) T-cells. *Retrovirology* 9, 87.

Fischer, E.S., Scrima, A., Böhm, K., Matsumoto, S., Lingaraju, G.M., Faty, M., Yasuda, T., Cavadini, S., Wakasugi, M., Hanaoka, F., et al. (2011). The molecular basis of CRL4DDB2/CSA ubiquitin ligase architecture, targeting, and activation. *Cell* 147, 1024–1039.

Franzolin, E., Pontarin, G., Rampazzo, C., Miazzi, C., Ferraro, P., Palumbo, E., Reichard, P., and Bianchi, V. (2013). The deoxynucleotide triphosphohydrolase SAMHD1 is a major regulator of DNA precursor pools in mammalian cells. *Proc. Natl. Acad. Sci. USA* 110, 14272–14277.

Fregoso, O.I., Ahn, J., Wang, C., Mehrens, J., Skowronski, J., and Emerman, M. (2013). Evolutionary toggling of Vpx/Vpr specificity results in divergent recognition of the restriction factor SAMHD1. *PLoS Pathog.* 9, e1003496.

Goldstone, D.C., Ennis-Adeniran, V., Hedden, J.J., Groom, H.C., Rice, G.I., Christodoulou, E., Walker, P.A., Kelly, G., Haire, L.F., Yap, M.W., et al. (2011). HIV-1 restriction factor SAMHD1 is a deoxynucleoside triphosphate triphosphohydrolase. *Nature* 480, 379–382.

Guo, Y., Dong, L., Qiu, X., Wang, Y., Zhang, B., Liu, H., Yu, Y., Zang, Y., Yang, M., and Huang, Z. (2014). Structural basis for hijacking CBF- $\beta$  and CUL5 E3 ligase complex by HIV-1 Vif. *Nature* 505, 229–233.

Hatzioannou, T., and Bieniasz, P.D. (2011). Antiretroviral restriction factors. *Curr. Opin. Virol.* 1, 526–532.

Hatzioannou, T., Del Prete, G.Q., Keele, B.F., Estes, J.D., McNatt, M.W., Bitzegeio, J., Raymond, A., Rodriguez, A., Schmidt, F., Mac Trubey, C., et al. (2014). HIV-1-induced AIDS in monkeys. *Science* 344, 1401–1405.

Hofmann, H., Logue, E.C., Bloch, N., Daddacha, W., Polsky, S.B., Schultz, M.L., Kim, B., and Landau, N.R. (2012). The Vpx lentiviral accessory protein targets SAMHD1 for degradation in the nucleus. *J. Virol.* 86, 12552–12560.

Hrecka, K., Gierszewska, M., Srivastava, S., Kozaczekiewicz, L., Swanson, S.K., Florens, L., Washburn, M.P., and Skowronski, J. (2007). Lentiviral Vpr usurps Cul4-DDB1[VprBP] E3 ubiquitin ligase to modulate cell cycle. *Proc. Natl. Acad. Sci. USA* 104, 11778–11783.

Hrecka, K., Hao, C., Gierszewska, M., Swanson, S.K., Kesik-Brodacka, M., Srivastava, S., Florens, L., Washburn, M.P., and Skowronski, J. (2011). Vpx relieves inhibition of HIV-1 infection of macrophages mediated by the SAMHD1 protein. *Nature* 474, 658–661.

Ji, X., Wu, Y., Yan, J., Mehrens, J., Yang, H., DeLucia, M., Hao, C., Gronenborn, A.M., Skowronski, J., Ahn, J., and Xiong, Y. (2013). Mechanism of allosteric activation of SAMHD1 by dGTP. *Nat. Struct. Mol. Biol.* 20, 1304–1309.

Karplus, P.A., and Diederichs, K. (2012). Linking crystallographic model and data quality. *Science* 336, 1030–1033.

Kim, B., Nguyen, L.A., Daddacha, W., and Hollenbaugh, J.A. (2012). Tight interplay among SAMHD1 protein level, cellular dNTP levels, and HIV-1 proviral DNA synthesis kinetics in human primary monocyte-derived macrophages. *J. Biol. Chem.* 287, 21570–21574.

Kretschmer, S., Wolf, C., König, N., Staroske, W., Guck, J., Häusler, M., Luksch, H., Nguyen, L.A., Kim, B., Alexopoulou, D., et al. (2015). SAMHD1 prevents autoimmunity by maintaining genome stability. *Ann. Rheum. Dis.* 74, e17.

Laguerre, N., Sobhian, B., Casartelli, N., Ringeard, M., Chable-Bessia, C., Ségéral, E., Yatim, A., Emiliani, S., Schwartz, O., and Benkirane, M. (2011). SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. *Nature* 474, 654–657.

Laguerre, N., Rahm, N., Sobhian, B., Chable-Bessia, C., Münch, J., Snoeck, J., Sauter, D., Switzer, W.M., Heneine, W., Kirchhoff, F., et al. (2012). Evolutionary



- and functional analyses of the interaction between the myeloid restriction factor SAMHD1 and the lentiviral Vpx protein. *Cell Host Microbe* 11, 205–217.
- Laguet, N., Br gnard, C., Hue, P., Basbous, J., Yatim, A., Larroque, M., Kirchhoff, F., Constantinou, A., Sobhian, B., and Benkirane, M. (2014). Premature activation of the SLX4 complex by Vpr promotes G2/M arrest and escape from innate immune sensing. *Cell* 156, 134–145.
- Lahouassa, H., Daddacha, W., Hofmann, H., Ayinde, D., Logue, E.C., Dragin, L., Bloch, N., Maudet, C., Bertrand, M., Gramberg, T., et al. (2012). SAMHD1 restricts the replication of human immunodeficiency virus type 1 by depleting the intracellular pool of deoxynucleoside triphosphates. *Nat. Immunol.* 13, 223–228.
- Le Rouzic, E., Belaidouni, N., Estrabaud, E., Morel, M., Rain, J.C., Transy, C., and Margottin-Goguet, F. (2007). HIV1 Vpr arrests the cell cycle by recruiting DCAF1/VprBP, a receptor of the Cul4-DDB1 ubiquitin ligase. *Cell Cycle* 6, 182–188.
- Lim, E.S., Fregoso, O.I., McCoy, C.O., Matsen, F.A., Malik, H.S., and Emerman, M. (2012). The ability of primate lentiviruses to degrade the monocyte restriction factor SAMHD1 preceded the birth of the viral accessory protein Vpx. *Cell Host Microbe* 11, 194–204.
- Malim, M.H., and Bieniasz, P.D. (2012). HIV Restriction Factors and Mechanisms of Evasion. *Cold Spring Harb. Perspect. Med.* 2, a006940.
- Miazzi, C., Ferraro, P., Pontarin, G., Rampazzo, C., Reichard, P., and Bianchi, V. (2014). Allosteric regulation of the human and mouse deoxyribonucleotide triphosphohydrolase sterile  $\alpha$ -motif/histidine-aspartate domain-containing protein 1 (SAMHD1). *J. Biol. Chem.* 289, 18339–18346.
- Mitchell, R.S., Katsura, C., Skasko, M.A., Fitzpatrick, K., Lau, D., Ruiz, A., Stephens, E.B., Margottin-Goguet, F., Benarous, R., and Guatelli, J.C. (2009). Vpu antagonizes BST-2-mediated restriction of HIV-1 release via beta-TrCP and endo-lysosomal trafficking. *PLoS Pathog.* 5, e1000450.
- Miyake, A., Fujita, M., Fujino, H., Koga, R., Kawamura, S., Otsuka, M., Ode, H., Iwatani, Y., Sakai, Y., Doi, N., et al. (2014). Poly-proline motif in HIV-2 Vpx is critical for its efficient translation. *J. Gen. Virol.* 95, 179–189.
- Neil, S.J., Zang, T., and Bieniasz, P.D. (2008). Tetherin inhibits retrovirus release and is antagonized by HIV-1 Vpu. *Nature* 451, 425–430.
- Pancio, H.A., Vander Heyden, N., and Ratner, L. (2000). The C-terminal proline-rich tail of human immunodeficiency virus type 2 Vpx is necessary for nuclear localization of the viral preintegration complex in nondividing cells. *J. Virol.* 74, 6162–6167.
- Pauls, E., Ruiz, A., Badia, R., Permanyer, M., Gubern, A., Riveira-Mu oz, E., Torres-Torronteras, J., Alvarez, M., Mothe, B., Brander, C., et al. (2014). Cell cycle control and HIV-1 susceptibility are linked by CDK6-dependent CDK2 phosphorylation of SAMHD1 in myeloid and lymphoid cells. *J. Immunol.* 193, 1988–1997.
- Powell, R.D., Holland, P.J., Hollis, T., and Perrino, F.W. (2011). Aicardi-Goutieres syndrome gene and HIV-1 restriction factor SAMHD1 is a dGTP-regulated deoxynucleotide triphosphohydrolase. *J. Biol. Chem.* 286, 43596–43600.
- Rehwinkel, J., Maelfait, J., Bridgeman, A., Rigby, R., Hayward, B., Liberatore, R.A., Bieniasz, P.D., Towers, G.J., Moita, L.F., Crow, Y.J., et al. (2013). SAMHD1-dependent retroviral control and escape in mice. *EMBO J.* 32, 2454–2462.
- Rice, G.I., Bond, J., Asipu, A., Brunette, R.L., Manfield, I.W., Carr, I.M., Fuller, J.C., Jackson, R.M., Lamb, T., Briggs, T.A., et al. (2009). Mutations involved in Aicardi-Goutieres syndrome implicate SAMHD1 as regulator of the innate immune response. *Nat. Genet.* 41, 829–832.
- Ryoo, J., Choi, J., Oh, C., Kim, S., Seo, M., Kim, S.Y., Seo, D., Kim, J., White, T.E., Brandariz-Nu ez, A., et al. (2014). The ribonuclease activity of SAMHD1 is required for HIV-1 restriction. *Nat. Med.* 20, 936–941.
- Schwefel, D., Groom, H.C., Boucherit, V.C., Christodoulou, E., Walker, P.A., Stoye, J.P., Bishop, K.N., and Taylor, I.A. (2014). Structural basis of lentiviral subversion of a cellular protein degradation pathway. *Nature* 505, 234–238.
- Sharp, P.M., and Hahn, B.H. (2011). Origins of HIV and the AIDS pandemic. *Cold Spring Harb. Perspect. Med.* 1, a006841.
- Sheehy, A.M., Gaddis, N.C., Choi, J.D., and Malim, M.H. (2002). Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* 418, 646–650.
- Spragg, C.J., and Emerman, M. (2013). Antagonism of SAMHD1 is actively maintained in natural infections of simian immunodeficiency virus. *Proc. Natl. Acad. Sci. USA* 110, 21136–21141.
- Srivastava, S., Swanson, S.K., Manel, N., Florens, L., Washburn, M.P., and Skowronski, J. (2008). Lentiviral Vpx accessory factor targets VprBP/DCAF1 substrate adaptor for cullin 4 E3 ubiquitin ligase to enable macrophage infection. *PLoS Pathog.* 4, e1000059.
- St Gelais, C., de Silva, S., Amie, S.M., Coleman, C.M., Hoy, H., Hollenbaugh, J.A., Kim, B., and Wu, L. (2012). SAMHD1 restricts HIV-1 infection in dendritic cells (DCs) by dNTP depletion, but its expression in DCs and primary CD4+ T-lymphocytes cannot be upregulated by interferons. *Retrovirology* 9, 105.
- Strebel, K. (2013). HIV accessory proteins versus host restriction factors. *Curr. Opin. Virol.* 3, 692–699.
- Stremelau, M., Owens, C.M., Perron, M.J., Kiessling, M., Autissier, P., and Sodroski, J. (2004). The cytoplasmic body component TRIM5alpha restricts HIV-1 infection in Old World monkeys. *Nature* 427, 848–853.
- Towers, G.J., and Noursadeghi, M. (2014). Interactions between HIV-1 and the cell-autonomous innate immune system. *Cell Host Microbe* 16, 10–18.
- T ngler, V., Staroske, W., Kind, B., Dobrick, M., Kretschmer, S., Schmidt, F., Krug, C., Lorenz, M., Chara, O., Schwillie, P., and Lee-Kirsch, M.A. (2013). Single-stranded nucleic acids promote SAMHD1 complex formation. *J. Mol. Med.* 91, 759–770.
- Wei, W., Guo, H., Gao, Q., Markham, R., and Yu, X.F. (2014). Variation of two primate lineage-specific residues in human SAMHD1 confers resistance to N terminus-targeted SIV Vpx proteins. *J. Virol.* 88, 583–591.
- White, T.E., Brandariz-Nu ez, A., Valle-Casuso, J.C., Amie, S., Nguyen, L.A., Kim, B., Tuzova, M., and Diaz-Griffero, F. (2013). The retroviral restriction ability of SAMHD1, but not its deoxynucleotide triphosphohydrolase activity, is regulated by phosphorylation. *Cell Host Microbe* 13, 441–451.
- Wolf, D., and Goff, S.P. (2008). Host restriction factors blocking retroviral replication. *Annu. Rev. Genet.* 42, 143–163.
- Zhou, Y., Lu, Y., and Ratner, L. (1998). Arginine residues in the C-terminus of HIV-1 Vpr are important for nuclear localization and cell cycle arrest. *Virology* 242, 414–424.
- Zhu, C., Gao, W., Zhao, K., Qin, X., Zhang, Y., Peng, X., Zhang, L., Dong, Y., Zhang, W., Li, P., et al. (2013). Structural insight into dGTP-dependent activation of tetrameric SAMHD1 deoxynucleoside triphosphate triphosphohydrolase. *Nat. Commun.* 4, 2722.
- Zimmerman, E.S., Schulman, B.A., and Zheng, N. (2010). Structural assembly of cullin-RING ubiquitin ligase complexes. *Curr. Opin. Struct. Biol.* 20, 714–721.